


KRT8/KRT18 monoclonal antibody, clone CK 209 (FITC)

Catalog # MAB12882 Size 250 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human liver with KRT8/KRT18 monoclonal antibody, clone CK 209 (FITC) (Cat # MAB12882).

Specification

Product Description	Mouse monoclonal antibody raised against human KRT8/KRT18.
Immunogen	Human breast cancer PMC42 cells.
Host	Mouse
Reactivity	Human
Form	Liquid
Conjugation	FITC
Purification	Protein A purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (2-10 ug/mL) Immunocytochemistry (10-20 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (1% BSA, 0.05% sodium azide).

Storage Instruction

Store at 4°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human liver with KRT8/KRT18 monoclonal antibody, clone CK 209 (FITC) (Cat # MAB12882).

- Immunocytochemistry
- Flow Cytometry

Gene Info — KRT8

Entrez GeneID[3856](#)**Gene Name**

KRT8

Gene Alias

CARD2, CK8, CYK8, K2C8, K8, KO

Gene Description

keratin 8

Omim ID[148060](#) [215600](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene is a member of the type II keratin family clustered on the long arm of chromosome 12. Type I and type II keratins heteropolymerize to form intermediate-sized filaments in the cytoplasm of epithelial cells. The product of this gene typically dimerizes with keratin 18 to form an intermediate filament in simple single-layered epithelial cells. This protein plays a role in maintaining cellular structural integrity and also functions in signal transduction and cellular differentiation. Mutations in this gene cause cryptogenic cirrhosis. [provided by RefSeq]

Other Designations

cytokeratin 8|keratin, type II cytoskeletal 8

Gene Info — KRT18

Entrez GeneID[3875](#)

Gene Name	KRT18
Gene Alias	CYK18, K18
Gene Description	keratin 18
Omim ID	148070 215600
Gene Ontology	Hyperlink
Gene Summary	KRT18 encodes the type I intermediate filament chain keratin 18. Keratin 18, together with its filament partner keratin 8, are perhaps the most commonly found members of the intermediate filament gene family. They are expressed in single layer epithelial tissues of the body. Mutations in this gene have been linked to cryptogenic cirrhosis. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq]
Other Designations	cell proliferation-inducing protein 46 cytokeratin 18

Pathway

- [Pathogenic Escherichia coli infection - EHEC](#)

Disease

- [Alzheimer disease](#)
- [Cerebral Amyloid Angiopathy](#)
- [Chronic Disease](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Disease Progression](#)
- [Drug-Induced Liver Injury](#)
- [Drug-Induced Liver Injury](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Inflammatory Bowel Diseases](#)

- [Liver Cirrhosis](#)
- [Liver Cirrhosis](#)
- [Liver Failure](#)
- [Liver Failure](#)
- [Neuroblastoma](#)
- [Pancreatitis](#)