KRT7 monoclonal antibody, clone SPM270

Catalog # MAB13100 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung squamous cell carcinoma with KRT7 monoclonal antibody, clone SPM270 (Cat # MAB13100).

Specification

Product Description	Mouse monoclonal antibody raised against native human KRT7.
Immunogen	OTN 11 ovarian carcinoma cell line.
Host	Mouse
Theoretical MW (kDa)	55
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1
Recommend Usage	Flow Cytometry (0.5-1 ug/10^6 cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at -20 to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung squamous cell carcinoma with KRT7 monoclonal antibody, clone SPM270 (Cat # MAB13100).

- Immunofluorescence

- Flow Cytometry

Gene Info — KRT7

Entrez GeneID	3855
Protein Accession#	P08729
Gene Name	KRT7
Gene Alias	CK7, K2C7, K7, MGC129731, MGC3625, SCL
Gene Description	keratin 7
Omim ID	148059
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coex pressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is s pecifically expressed in the simple epithelia lining the cavities of the internal organs and in the gla nd ducts and blood vessels. The genes encoding the type II cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing may result in several transcript variants; however, not all variants have been fully described. [provided by RefSeq
Other Designations	cytokeratin 7|keratin, 55K type II cytoskeletal|keratin, simple epithelial type I, K7|keratin, type II cyto skeletal 7|sarcolectin|type II mesothelial keratin K7

Copyright © 2023 Abnova Corporation. All Rights Reserved.