


CDK1 monoclonal antibody, clone CDK1/873

Catalog # MAB13226 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil using CDK1 monoclonal antibody, clone CDK1/873 (Cat # MAB13226).

Specification	
Product Description	Mouse monoclonal antibody raised against full length recombinant human CDK1.
Immunogen	Recombinant protein corresponding to full length human CDK1.
Host	Mouse
Theoretical MW (kDa)	34
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	lgG2a, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/million cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (2-4 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.



Storage Instruction

Store at -20 to -80°C.

Aliquot to avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
 Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil using CDK1 monoclonal antibody, clone CDK1/873 (Cat # MAB13226).
- Immunofluorescence
- Flow Cytometry

Gene Info — CDC2	
Entrez GenelD	983
Protein Accession#	<u>P06493</u>
Gene Name	CDC2
Gene Alias	CDC28A, CDK1, DKFZp686L20222, MGC111195
Gene Description	cell division cycle 2, G1 to S and G2 to M
Omim ID	<u>116940</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting f actor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitot ic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phos phorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle
	control. Alternatively spliced transcript variants encoding different isoforms have been found for thi s gene. [provided by RefSeq

Pathway



- Cell cycle
- Gap junction
- p53 signaling pathway

Disease

- Alzheimer disease
- Breast cancer
- Breast Neoplasms
- Dementia
- Genetic Predisposition to Disease
- Lung Neoplasms
- Pulmonary Disease