


CD59 monoclonal antibody, clone VJ1/12,2 (FITC)

Catalog # MAB13954 Size 5 x 100 reactions

Applications



Flow Cytometry

Flow cytometric analysis of human normal whole blood with CD59 monoclonal antibody, clone VJ1/12,2 (FITC) (Cat # MAB13954).

| Specification | |
|----------------------|--|
| Product Description | Mouse monoclonal antibody raised against human CD59. |
| Immunogen | TNF activated HUVEC cells. |
| Host | Mouse |
| Theoretical MW (kDa) | 18-20 |
| Reactivity | Human |
| Form | Liquid |
| Conjugation | FΠC |
| Purification | Protein A/G purification |
| Purity | >90% |
| Isotype | lgG2a |



Product Information

| Recommend Usage | Flow Cytometry (20 uL/10 ⁶ cells) The optimal working dilution should be determined by the end user. |
|---------------------|---|
| Storage Buffer | In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide). |
| Storage Instruction | Store in the dark at 4°C. Avoid prolonged exposure to light. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only. |

Applications

Flow Cytometry

Flow cytometric analysis of human normal whole blood with CD59 monoclonal antibody, clone VJ1/12,2 (FITC) (Cat # MAB13954).

| Gene Info — CD59 | |
|--------------------|---|
| Entrez GeneID | <u>966</u> |
| Protein Accession# | <u>P13987</u> |
| Gene Name | CD59 |
| Gene Alias | 16.3A5, 1F5, EJ16, EJ30, EL32, FLJ38134, FLJ92039, G344, HRF-20, HRF20, MAC-IP, MACI F, MEM43, MGC2354, MIC11, MIN1, MIN2, MIN3, MIRL, MSK21, p18-20 |
| Gene Description | CD59 molecule, complement regulatory protein |
| Omim ID | 107271 |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | This gene encodes a cell surface glycoprotein that regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. This protein also plays a role in signal transduction pathways in the activation of T cells. Mutations in this gene cause CD59 deficiency, a disease resulting in hemolytic anemia and thrombosis, and which causes cerebral infarction. Multiple alternatively spliced transcript variants, which encode the same protein, have been identified for this gene. [provided by RefSeq |
| Other Designations | 20 kDa homologous restriction factor CD59 antigen CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344) CD59 glycoprotein Ly-6-like protein T cell-activating protein human leukocyte antigen MIC11 lymphocytic a |



Pathway

- Complement and coagulation cascades
- Hematopoietic cell lineage

Disease

- Genetic Predisposition to Disease
- Lymphoma
- Macular Degeneration