


MYC monoclonal antibody, clone MYC909

Catalog # MAB14371 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human cervical carcinoma with MYC monoclonal antibody, clone MYC909 (Cat # MAB14371).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human MYC.
Immunogen	Recombinant protein corresponding to full length human MYC.
Host	Mouse
Reactivity	Human
Specificity	This monoclonal antibody shows no cross-reaction with v-myc.
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% BSA, 0.05% sodium azide).

Storage Instruction

Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human cervical carcinoma with MYC monoclonal antibody, clone MYC909 (Cat # MAB14371).

- Immunofluorescence

- Flow Cytometry

Gene Info — MYC

Entrez GeneID[4609](#)**Protein Accession#**[P01106](#)**Gene Name**

MYC

Gene Alias

bHLHe39, c-Myc

Gene Description

v-myc myelocytomatosis viral oncogene homolog (avian)

Omim ID[113970 190080](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemia and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq]

Other Designations

avian myelocytomatosis viral oncogene homolog|myc proto-oncogene protein|v-myc avian myelocytomatosis viral oncogene homolog

Pathway

- [Acute myeloid leukemia](#)
- [Bladder cancer](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [TGF-beta signaling pathway](#)
- [Thyroid cancer](#)
- [Wnt signaling pathway](#)

Disease

- [Alzheimer disease](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chromosome Aberrations](#)
- [Colorectal Neoplasms](#)
- [Diabetes Complications](#)
- [Disease Progression](#)

- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Mouth Neoplasms](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Thyroid Diseases](#)
- [Thyroid Neoplasms](#)
- [Translocation](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)