


NAPSA monoclonal antibody, clone NAPSA/1238

Catalog # MAB14422 Size 100 ug


Applications



Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis with NAPSA monoclonal antibody, clone NAPSA/1238 (Cat # MAB14422):

1. K562 cell line
2. HEK293 cell line
3. A549 cell line



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung adenocarcinoma with NAPSA monoclonal antibody, clone NAPSA/1238 (Cat # MAB14422).

Specification

| | |
|----------------------------|---|
| Product Description | Mouse monoclonal antibody raised against partial recombinant human NAPSA. |
| Immunogen | Recombinant protein corresponding to human NAPSA. |
| Host | Mouse |
| Reactivity | Human |
| Form | Liquid |
| Purification | Protein A purification |

| | |
|---------------------|---|
| Isotype | IgG, kappa |
| Recommend Usage | Flow Cytometry (0.5-1 ug/million cells) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) Western Blotting (1-2 ug/mL) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In 10 mM PBS (0.05% BSA and 0.05% azide). |
| Storage Instruction | Store at 4°C. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Applications

- Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis with NAPSA monoclonal antibody, clone NAPSA/1238 (Cat # MAB14422):

1. K562 cell line
2. HEK293 cell line
3. A549 cell line

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung adenocarcinoma with NAPSA monoclonal antibody, clone NAPSA/1238 (Cat # MAB14422).

- Immunofluorescence

- Flow Cytometry

Gene Info — NAPSA

| | |
|--------------------|------------------------------|
| Entrez GeneID | 9476 |
| Protein Accession# | O96009 |
| Gene Name | NAPSA |
| Gene Alias | KAP, Kdap, NAP1, NAPA, SNAPA |
| Gene Description | napsin A aspartic peptidase |
| Omim ID | 605631 |

Gene Ontology

[Hyperlink](#)

Gene Summary

The activation peptides of aspartic proteinases plays role as inhibitors of the active site. These peptide segments, or pro-parts, are deemed important for correct folding, targeting, and control of the activation of aspartic proteinase zymogens. The pronapsin A gene is expressed predominantly in lung and kidney. Its translation product is predicted to be a fully functional, glycosylated aspartic proteinase precursor containing an RGD motif and an additional 18 residues at its C-terminus. [provided by RefSeq]

Other Designations

napsin A|pronapsin A

Publication Reference

- [Napsin A expression in lung and kidney neoplasia: a review and update.](#)

Nelson G Ordonez.

Advances in Anatomic Pathology 2012 Jan; 19(1):66.

Application: IHC-P, Human, Human lung adenocarcinomas, Human renal cell carcinomas

- [Combination of napsin A and TTF-1 immunohistochemistry helps in differentiating primary lung adenocarcinoma from metastatic carcinoma in the lung.](#)

Jiqing Ye, Jennifer J Findeis-Hosey, Qi Yang, Lorelee A McMahon, Jorge L Yao, Faqian Li, Haodong Xu.

Applied Immunohistochemistry & Molecular Morphology 2011 Jul; 19(4):313.

Application: IHC-P, Human, Human lung adenocarcinoma, Human renal cell carcinomas, Human tissue microarray

- [Napsin A and thyroid transcription factor-1 expression in carcinomas of the lung, breast, pancreas, colon, kidney, thyroid, and malignant mesothelioma.](#)

Justin A Bishop, Rajni Sharma, Peter B Illei.

Human Pathology 2010 Jan; 41(1):20.

Application: IHC-P, Human, Human tissue microarray

Pathway

- [Lysosome](#)