


# CD3E monoclonal antibody, clone PC3/188A

Catalog # MAB14699      Size 100 ug

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with CD3E monoclonal antibody, clone PC3/188A (Cat # MAB14699).

## Specification

<b>Product Description</b>	Mouse monoclonal antibody raised against synthetic peptide of human CD3E.
<b>Immunogen</b>	A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic domain of human CD3E.
<b>Host</b>	Mouse
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Purification</b>	Protein A purification
<b>Isotype</b>	IgG1
<b>Recommend Usage</b>	Flow Cytometry (0.5-1 ug/million cells) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) Western Blot (1-2 ug/mL) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	In 10 mM PBS (0.05% BSA and 0.05% azide).

**Storage Instruction**

Store at 4°C.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)  
Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with CD3E monoclonal antibody, clone PC3/188A (Cat # MAB14699).
- Immunofluorescence
- Flow Cytometry

## Gene Info — CD3E

Entrez GenelD	<a href="#">916</a>
Gene Name	CD3E
Gene Alias	FLJ18683, T3E, TCRE
Gene Description	CD3e molecule, epsilon (CD3-TCR complex)
Omim ID	<a href="#">186830</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women. [provided by RefSeq]
Other Designations	CD3-epsilon CD3E antigen, epsilon polypeptide CD3e antigen, epsilon polypeptide (TiT3 complex) T-cell antigen receptor complex, epsilon subunit of T3 T-cell surface antigen T3/Leu-4 epsilon chain T-cell surface glycoprotein CD3 epsilon chain

## Publication Reference

- [The expression of the CD3 antigen in Hodgkin's disease.](#)

Cibull ML, Stein H, Gatter KC, Mason DY.

Histopathology 1989 Dec; 15(6):599.

- [Detection of T cells in paraffin wax embedded tissue using antibodies against a peptide sequence from the CD3 antigen.](#)

D Y Mason, J Cordell, M Brown, G Pallesen, E Ralfkiaer, J Rothbard, M Crumpton, K C Gatter.

Journal of Clinical Pathology 1989 Nov; 42(11):1194.

Application: IHC-Fr, IHC-P, IP, RIA, Human, Human T cell lymphoma

## Pathway

- [Hematopoietic cell lineage](#)
- [Primary immunodeficiency](#)
- [T cell receptor signaling pathway](#)

## Disease

- [Asthma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Inflammation](#)