


TOP2A monoclonal antibody, clone TOP2A/1362

Catalog # MAB15015 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil (A) and human bladder carcinoma (B) with TOP2A monoclonal antibody, clone TOP2A/1362 (Cat # MAB15015).

Specification

Product Description	Mouse monoclonal antibody raised against partial recombinant human TOP2A.
Immunogen	Recombinant protein corresponding to amino acids 1352-1493 of human TOP2A.
Host	Mouse
Theoretical MW (kDa)	170
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG2b, kappa

Recommend Usage

Flow Cytometry (0.5-1 ug/10⁶ cells)
Immunofluorescence (1-2 ug/mL)
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL)
Western Blotting (0.5-1 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer

In 10 mM PBS.

Storage Instruction

Store at -20 to -80°C.
Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil (A) and human bladder carcinoma (B) with TOP2A monoclonal antibody, clone TOP2A/1362 (Cat # MAB15015).
- Immunofluorescence
- Flow Cytometry

Gene Info — TOP2A

Entrez GeneID [7153](#)

Protein Accession# [P11388](#)

Gene Name TOP2A

Gene Alias TOP2, TP2A

Gene Description topoisomerase (DNA) II alpha 170kDa

Omim ID [126430](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic state of DNA during transcription. This nuclear enzyme is involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. It catalyzes the transient breaking and rejoining of two strands of duplex DNA which allows the strands to pass through one another, thus altering the topology of DNA. Two forms of this enzyme exist as likely products of a gene duplication event. The gene encoding this form, alpha, is localized to chromosome 17 and the beta gene is localized to chromosome 3. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also play a role in ataxia-telangiectasia. [provided by RefSeq]

Other Designations

DNA topoisomerase II, 170 kD|DNA topoisomerase II, alpha isozyme|topoisomerase (DNA) II alpha (170kD)

Publication Reference

- [Characterization and immunological identification of cDNA clones encoding two human DNA topoisomerase II isozymes.](#)
Chung TD, Drake FH, Tan KB, Per SR, Crooke ST, Mirabelli CK.
PNAS 1989 Dec; 86(23):9431.

Disease

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Leukemia](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Recurrence](#)
- [Stomach Neoplasms](#)