
RecomAb™

CDKN1B monoclonal antibody, clone RM302


Catalog # MAB16928 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human breast cancer.



Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis of MCF-7 cell lysates.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CDKN1B.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to C-terminus of human CDKN 1B.
Reactivity	Human, Mouse
Specificity	This antibody reacts to human p27Kip1 (Cyclin-dependent kinase inhibitor 1B). It may also react to m ouse p27Kip1, as predicted by immunogen homology.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:100-200) Western Blot (1:500-1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining of human breast cancer.

- Western Blot (Cell lysate)

  Western Blot (Cell lysate) analysis of MCF-7 cell lysates.

Gene Info — CDKN1B

Entrez GeneID	1027
Gene Name	CDKN1B
Gene Alias	CDKN4, KIP1, MEN1B, MEN4, P27KIP1
Gene Description	cyclin-dependent kinase inhibitor 1B (p27, Kip1)
Omim ID	600778 610755
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CD K inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-C DK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degrad ation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubi quitination by SCF complexes, is required for the cellular transition from quiescence to the prolifer ative state. [provided by RefSeq

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Other Designations	cyclin-dependent kinase inhibitor 1B

Pathway

- Cell cycle
- Chronic myeloid leukemia
- ErbB signaling pathway
- Pathways in cancer
- Prostate cancer
- Small cell lung cancer

Disease

- Acromegaly
- Adenocarcinoma
- Alzheimer disease
- Ataxia telangiectasia
- Atherosclerosis
- Breast cancer
- Breast Neoplasms
- Carcinoma
- Cardiovascular Diseases
- Chromosome Aberrations
- Cognition
- Colon cancer
- Colorectal Neoplasms
- Coronary Artery Disease

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Coronary Restenosis
- Diabetes Mellitus
- Disease Progression
- Edema
- Endometriosis
- Esophageal Neoplasms
- Genetic Predisposition to Disease
- Graft Occlusion
- Head and Neck Neoplasms
- Hyperparathyroidism
- Kidney Failure
- Leukemia
- Liver Neoplasms
- Lung Neoplasms
- Lymphatic Metastasis
- Lymphoma
- Mouth Neoplasms
- Multiple endocrine neoplasia
- Multiple Endocrine Neoplasia Type 1
- Myocardial Infarction
- Neoplasm Invasiveness
- Neoplasm Metastasis
- Neoplasm Recurrence
- Neoplasms
- Ovarian cancer
- Ovarian Neoplasms

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Pancreatic cancer
- Pancreatic Neoplasms
- Parathyroid Neoplasms
- Precancerous Conditions
- Primary Ovarian Insufficiency
- Prostate cancer
- Prostatic Neoplasms
- Pulmonary Disease
- Stomach Neoplasms
- Thyroid Neoplasms
- Urinary Bladder Neoplasms
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.