


HIST2H4A(20Me3) monoclonal antibody, clone 7A2E10


Catalog # MAB17145 Size 100 ug

Applications



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of HIST2H4A(20Me3) monoclonal antibody, clone 7A2E10.



Flow Cytometry

Flow cytometric analysis of Raji cells with HIST2H4A(20Me3) monoclonal antibody (green) and negative control (red).

Specification

Product Description	Mouse monoclonal antibody raised against synthetic peptide of human HIST2H4A(20Me3).
Immunogen	A synthetic peptide corresponding to amino acid of human HIST2H4A(20Me3).
Host	Mouse
Theoretical MW (kDa)	11.4kDa
Reactivity	Human
Form	Liquid

Isotype	IgG1
Recommend Usage	ELISA (1:10000) Western Blot Immunohistochemistry Immunocytochemistry Flow Cytometry (1:200-1:400) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis of HIST2H4A(20Me3) monoclonal antibody, clone 7A2E10.

- Flow Cytometry

Flow cytometric analysis of Raji cells with HIST2H4A(20Me3) monoclonal antibody (green) and negative control (red).

Gene Info — HIST2H4A

Entrez GeneID	8370
Gene Name	HIST2H4A
Gene Alias	FO108, H4, H4/n, H4F2, H4FN, HIST2H4
Gene Description	histone cluster 2, H4a
Omim ID	142750
Gene Ontology	Hyperlink

Gene Summary

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy. [provided by RefSeq]

Other Designations

H4 histone family, member N|H4 histone, family 2|OTTHUMP00000013906|OTTHUMP00000194862|OTTHUMP00000194863|histone 2, H4a|histone IV, family 2

Pathway

- [Systemic lupus erythematosus](#)