


IL2RB monoclonal antibody, clone 1G11H3


Catalog # MAB17626 Size 100 ug

Applications




Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) IL2RB-hlgGfc transfected HEK293 cell lysate.



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of IL2RB monoclonal antibody, clone 1G11H3.



Flow Cytometry

Flow cytometric analysis of Ramos cells with IL2RB monoclonal antibody (green) and negative control (red).

Specification

Product Description	Mouse monoclonal antibody raised against recombinant human IL2RB.
Immunogen	Recombinant protein corresponding to amino acid 27-240 of human IL2RB from <i>E. coli</i> .
Host	Mouse
Theoretical MW (kDa)	61
Reactivity	Human
Form	Liquid
Isotype	IgG2b
Recommend Usage	ELISA (1:10000) Western Blot (1:500-1:2000) Immunocytochemistry Flow Cytometry (1:200-1:400) Immunohistochemistry The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) IL2RB-hlgGfc transfected HEK293 cell lysate.

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis of IL2RB monoclonal antibody, clone 1G11H3.

- Flow Cytometry

Flow cytometric analysis of Ramos cells with IL2RB monoclonal antibody (green) and negative control (red).

Gene Info — IL2RB

Entrez GeneID [3560](#)

Gene Name IL2RB

Gene Alias	CD122, P70-75
Gene Description	interleukin 2 receptor, beta
Omim ID	146710
Gene Ontology	Hyperlink
Gene Summary	The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is present in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha s ubunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha /beta subunit heterodimer, while the high affinity form consists of an alpha/beta/gamma subunit he terotrimer. Both the intermediate and high affinity forms of the receptor are involved in receptor-m ediated endocytosis and transduction of mitogenic signals from interleukin 2. The protein encode d by this gene represents the beta subunit and is a type I membrane protein. [provided by RefSeq
Other Designations	CD122 antigen OTTHUMP00000028799 high affinity IL-2 receptor beta subunit interleukin 2 rece ptor beta

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bipolar Disorder](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Chagas Disease](#)

- [Colitis](#)
- [Connective Tissue Diseases](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometriosis](#)
- [Esophagitis](#)
- [Fetal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Hyperparathyroidism](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leishmaniasis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Measles](#)
- [Multiple Sclerosis](#)
- [Mumps](#)
- [Musculoskeletal Diseases](#)
- [Myasthenia Gravis](#)
- [Occupational Diseases](#)

- [Pain](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Radiation Injuries](#)
- [Radiation Pneumonitis](#)
- [Respiratory Syncytial Virus Infections](#)
- [Rubella](#)
- [Schizophrenia](#)
- [Skin Diseases](#)
- [Tobacco Use Disorder](#)
- [Viremia](#)
- [Wegener Granulomatosis](#)