


BCR monoclonal antibody, clone 6B5E12


Catalog # MAB17934 Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of Lane 1: HEK293 and Lane 2: BCR-hlgFc transfected HEK293 cell lysates with BCR monoclonal antibody, clone 6B5E12 (Cat # MAB17934).



Enzyme-linked Immunoabsorbent Assay

ELISA analysis with BCR monoclonal antibody, clone 6B5E12 (Cat # MAB17934).

Specification

Product Description	Mouse monoclonal antibody raised against partial recombinant human BCR.
Immunogen	Recombinant protein corresponding to amino acids 139-280 of human BCR.
Host	Mouse
Theoretical MW (kDa)	142.8
Reactivity	Human
Form	Liquid

Isotype	IgG1
Recommend Usage	ELISA (1:10000) Western Blot (1:100-1:500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of Lane 1: HEK293 and Lane 2: BCR-hlgFc transfected HEK293 cell lysates with BCR monoclonal antibody, clone 6B5E12 (Cat # MAB17934).

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis with BCR monoclonal antibody, clone 6B5E12 (Cat # MAB17934).

Gene Info — BCR

Entrez GeneID	613
Protein Accession#	P11274
Gene Name	BCR
Gene Alias	ALL, BCR-ABL1, BCR1, CML, D22S11, D22S662, FLJ16453, PHL
Gene Description	breakpoint cluster region
Omim ID	151410 608232
Gene Ontology	Hyperlink

Gene Summary

A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

bcr-abl1 e14a3 chimeric protein

Pathway

- [Chronic myeloid leukemia](#)
- [Pathways in cancer](#)

Disease

- [Alzheimer disease](#)
- [Angina Pectoris](#)
- [Bipolar Disorder](#)
- [Brief Psychiatric Rating Scale](#)
- [Chromosome Breakage](#)
- [Coronary Vasospasm](#)
- [Depressive Disorder](#)
- [Genetic Predisposition to Disease](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Pulmonary Disease](#)
- [Small Cell Lung Carcinoma](#)
- [Tobacco Use Disorder](#)
- [Urinary Bladder Neoplasms](#)

- [Werner syndrome](#)