


MAP2K3/MAP2K6 (phospho S189/S207) monoclonal antibody, clone D3

Catalog # MAB18961 Size 200 uL

Applications



Flow Cytometry

Flow cytometric analysis of HEK293T cells secondary antibody only negative control (blue) or treated with K252a (red) or treated with UV+TPA (green) using MAP2K3/MAP2K6 (phospho S189/S207) monoclonal antibody.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human MAP2K3/MAP2K6.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding S189/S207 of human MAP2K3/MAP2K6.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Protein A/G Purification
Isotype	IgG1k
Recommend Usage	Flow Cytometry (1 ug/mL - 0.001 ug/mL) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, pH 7.4 (50% glycerol, 0.02% sodium azide, 0.1% BSA).
Storage Instruction	Store at -20°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of HEK293T cells secondary antibody only negative control (blue) or treated with K252a (red) or treated with UV+TPA (green) using MAP2K3/MAP2K6 (phospho S189/S207) monoclonal antibody.

Gene Info — MAP2K3

Entrez GeneID	5606
Gene Name	MAP2K3
Gene Alias	MAPKK3, MEK3, MKK3, PRKMK3
Gene Description	mitogen-activated protein kinase kinase 3
Omim ID	602315
Gene Ontology	Hyperlink

Gene Summary	The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersinia pseudotuberculosis. Multiple alternatively spliced transcript variants that encode distinct isoforms have been reported for this gene. [provided by RefSeq]
--------------	---

Other Designations	MAP kinase kinase 3 MAPK/ERK kinase 3 OTTHUMP00000166044 dual specificity mitogen activated protein kinase kinase 3
--------------------	---

Gene Info — MAP2K6

Entrez GeneID	5608
---------------	----------------------

Gene Name	MAP2K6
Gene Alias	MAPKK6, MEK6, MKK6, PRKMK6, SAPKK3
Gene Description	mitogen-activated protein kinase kinase 6
Omim ID	601254
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis. [provided by RefSeq]
Other Designations	protein kinase, mitogen-activated, kinase 6 (MAP kinase kinase 6)

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [GnRH signaling pathway](#)
- [GnRH signaling pathway](#)
- [MAPK signaling pathway](#)
- [MAPK signaling pathway](#)
- [Toll-like receptor signaling pathway](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)

- [Edema](#)
- [Huntington disease](#)