


STAT3 (phospho Y705) monoclonal antibody, clone B12 (FITC)

Catalog # MAB19128 Size 10 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of Jurkat cells with STAT3 (phospho Y705) monoclonal antibody, clone B12 (FITC) (Cat # MAB19128). Secondary antibody only negative control (blue) or untreated (red) or treated with IFN α , IL-4 and pervanadate (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human STAT3.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y705 of human STAT3.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Conjugation	FITC
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (5 μ L/ 10^6 cells) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).

Storage Instruction

Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of Jurkat cells with STAT3 (phospho Y705) monoclonal antibody, clone B12 (FITC) (Cat # MAB19128). Secondary antibody only negative control (blue) or untreated (red) or treated with IFN α , IL-4 and pervanadate (green).

Gene Info — STAT3

Entrez GeneID

[6774](#)

Gene Name

STAT3

Gene Alias

APRF, FLJ20882, HIES, MGC16063

Gene Description

signal transducer and activator of transcription 3 (acute-phase response factor)

Omim ID

[102582](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations

DNA-binding protein APRF|acute-phase response factor|signal transducer and activator of transcription 3

Pathway

- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)

- [Chemokine signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)

Disease

- [Abortion](#)
- [Asthma](#)
- [Autoimmune Diseases](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Depressive Disorder](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [DNA Damage](#)
- [Eczema](#)

- [Edema](#)
- [Fatty Liver](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hepatitis C](#)
- [Infant](#)
- [Inflammation](#)
- [Inflammatory Bowel Diseases](#)
- [Insulin Resistance](#)
- [Kidney Neoplasms](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neoplasm Metastasis](#)
- [Obesity](#)
- [Ovarian Neoplasms](#)
- [Pulmonary Disease](#)
- [Rectal Fistula](#)
- [Respiratory Syncytial Virus Infections](#)
- [Spondylitis](#)

- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tooth Abnormalities](#)