


STAT6 (phospho Y641) monoclonal antibody, clone G12 (SureLight 488)

Catalog # MAB19158 Size 10 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of U937 cells with STAT6 (phospho Y641) monoclonal antibody, clone G12 (SureLight 488) (Cat # MAB19158). Untreated cells stained with secondary antibody as negative control (blue) or untreated (red) or treated with IFN α and IL-4 (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human STAT6.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y641 of human STAT6.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Conjugation	SureLight 488
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (5 μ L/ 10^6 cells) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).
Storage Instruction	Store at 4°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of U937 cells with STAT6 (phospho Y641) monoclonal antibody, clone G12 (SureLight 488) (Cat # MAB19158). Untreated cells stained with secondary antibody as negative control (blue) or untreated (red) or treated with IFN α and IL-4 (green).

Gene Info — STAT6

Entrez GeneID	6778
Gene Name	STAT6
Gene Alias	D12S1644, IL-4-STAT, STAT6B, STAT6C
Gene Description	signal transducer and activator of transcription 6, interleukin-4 induced
Omim ID	601512
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. [provided by RefSeq]
Other Designations	STAT, interleukin4-induced signal transducer and activator of transcription 6 transcription factor IL-4 STAT

Pathway

- [Jak-STAT signaling pathway](#)

Disease

- [Anaphylaxis](#)
- [Arthritis](#)
- [Ascariasis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Confusion](#)
- [Crohn Disease](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Hypersensitivity](#)
- [Edema](#)
- [Endometriosis](#)
- [Food Hypersensitivity](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Graves Disease](#)

- [Hypersensitivity](#)
- [Infant](#)
- [Infertility](#)
- [Inflammatory Bowel Diseases](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Diseases](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Nausea](#)
- [Neoplasms](#)
- [Nephrosis](#)
- [Nephrotic Syndrome](#)
- [Nut Hypersensitivity](#)
- [Pain](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)