HIST1H4A (mono methyl K16) monoclonal antibody, clone DEF-8

Catalog # MAB20163 Size 100 uL

Applications



Western Blot

Western Blot analysis of (1) NIH/3T3, (2) A549 cell lysate.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human HIST1H4A (mono methyl K16).
Immunogen	A synthetic peptide corresponding to human HIST1H4A (mono methyl K16).
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunocytochemistry (1:500-2000) Immunofluorescence (1:500-2000) Western Blot (1:500-2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and st ored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunofluorescence
- Immunocytochemistry
- Western Blot

  Western Blot analysis of (1) NIH/3T3, (2) A549 cell lysate.

Gene Info — HIST1H4A

Entrez GeneID	8359
Protein Accession#	P62805
Gene Name	HIST1H4A
Gene Alias	H4/a, H4FA
Gene Description	histone cluster 1, H4a
Omim ID	602822
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, an d H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and f unctions in the compaction of chromatin into higher order structures. This gene is intronless and e ncodes a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq
Other Designations	H4 histone family, member A|histone 1, H4a

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Systemic lupus erythematosus

Copyright © 2023 Abnova Corporation. All Rights Reserved.