


XRCC6 monoclonal antibody, clone AAIG-24

Catalog # MAB20430 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of HeLa cell lysate with XRCC6 monoclonal antibody, clone AAIG-24 (Cat # MAB20430).

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human XRCC6.

Immunogen A synthetic peptide corresponding to human XRCC6.

Host Rabbit

Theoretical MW (kDa) 69.843

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage

- Flow Cytometry (1:50)
- Immunocytochemistry (1:50-1:200)
- Immunofluorescence (1:50-1:200)
- Immunohistochemistry (1:50-1:200)
- Immunoprecipitation (1:50)
- Western Blot (1:500-1:2000)

The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of HeLa cell lysate with XRCC6 monoclonal antibody, clone AAIG-24 (Cat # MAB20430).

- Immunohistochemistry

- Immunocytochemistry

- Immunofluorescence

- Immunoprecipitation

- Flow Cytometry

Gene Info — XRCC6

Entrez GeneID	2547
Protein Accession#	P12956
Gene Name	XRCC6
Gene Alias	CTC75, CTCBF, G22P1, KU70, ML8, TLAA
Gene Description	X-ray repair complementing defective repair in Chinese hamster cells 6
Omim ID	152690
Gene Ontology	Hyperlink

Gene Summary

The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq]

Other Designations

ATP-dependent DNA helicase II, 70 kDa subunit|CTC box binding factor 75 kDa subunit|Ku autoantigen p70 subunit|Ku autoantigen, 70kDa|OTTHUMP00000028581|thyroid autoantigen 70kD (Ku antigen)|thyroid autoantigen 70kDa (Ku antigen)|thyroid-lupus autoantigen

Pathway

- [Non-homologous end-joining](#)

Disease

- [Adenocarcinoma](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Central Nervous System Neoplasms](#)
- [Diabetes Mellitus](#)
- [DNA Damage](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)

- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Male Urogenital Diseases](#)
- [Micronuclei](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pterygium](#)
- [Radiodermatitis](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)