


TRAF6 monoclonal antibody, clone BOF-20

Catalog # MAB20803 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human colon carcinoma with TRAF6 monoclonal antibody, clone BOF-20 (Cat # MAB20803).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human TRAF6.
Immunogen	A synthetic peptide corresponding to human TRAF6.
Host	Rabbit
Theoretical MW (kDa)	59.573
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Flow Cytometry (1:50-1:200) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-1:200) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human colon carcinoma with TRAF6 monoclonal antibody, clone BOF-20 (Cat # MAB20803).
- Flow Cytometry

Gene Info — TRAF6

Entrez GeneID [7189](#)

Protein Accession# [Q9Y4K3](#)

Gene Name TRAF6

Gene Alias MGC:3310, RNF85

Gene Description TNF receptor-associated factor 6

Omim ID [602355](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins are associated with, and mediate signal transduction from members of the TNF receptor superfamily. This protein mediates the signaling not only from the members of the TNF receptor superfamily, but also from the members of the Toll/IL-1 family. Signals from receptors such as CD40, TNFSF11/RANCE and IL-1 have been shown to be mediated by this protein. This protein also interacts with various protein kinases including IRAK1/IRAK, SRC and PKCzeta, which provides a link between distinct signaling pathways. This protein functions as a signal transducer in the NF-kappaB pathway that activates IkkappaB kinase (IKK) in response to proinflammatory cytokines. The interaction of this protein with UBE2N/UBC13, and UBE2V1/UEV1A, which are ubiquitin conjugating enzymes catalyzing the formation of polyubiquitin chains, has been found to be required for IKK activation by this protein. Two alternatively spliced transcript variants encoding identical proteins have been reported. [provided by RefSeq]

Other Designations

-

Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [Toll-like receptor signaling pathway](#)
- [Ubiquitin mediated proteolysis](#)

Disease

- [Alzheimer disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)

- [Hepatitis C](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Hypersensitivity](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Narcolepsy](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Thyroid Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)