


CDKN1B monoclonal antibody, clone DCS-72.F6 + KIP1/769

Catalog # MAB20867

Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human colon carcinoma with CDKN1B monoclonal antibody, clone DCS-72.F6 + KIP1/769 (Cat # MAB20867).

Specification

Product Description Mouse monoclonal antibody raised against human CDKN1B.

Immunogen Recombinant protein corresponding to human p27 and CDKN1B.

Host Mouse

Theoretical MW (kDa) 25-26

Reactivity Human

Form Liquid

Purification Protein A/G purification

Isotype IgG1, kappa

Recommend Usage

- Flow Cytometry (0.5-1 ug/10⁶ cells in 0.1 mL)
- Immunofluorescence (0.5-1 ug/mL)
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.25-0.5 ug/mL)
- Western Blot (0.5-1 ug/mL)
- The optimal working dilution should be determined by the end user.

Storage Buffer In 10 mM PBS (0.05% BSA, 0.05% sodium azide).

Storage Instruction

Store at 2 to 8°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human colon carcinoma with CDKN1B monoclonal antibody, clone DCS-72.F6 + KIP1/769 (Cat # MAB20867).

- Immunofluorescence
- Flow Cytometry

Gene Info — CDKN1B

Entrez GeneID [1027](#)**Protein Accession#** [P46527](#)**Gene Name** CDKN1B**Gene Alias** CDKN4, KIP1, MEN1B, MEN4, P27KIP1**Gene Description** cyclin-dependent kinase inhibitor 1B (p27, Kip1)**Omim ID** [600778](#) [610755](#)**Gene Ontology** [Hyperlink](#)

Gene Summary This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. [provided by RefSeq]

Other Designations cyclin-dependent kinase inhibitor 1B

Pathway

- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [ErbB signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)

Disease

- [Acromegaly](#)
- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Ataxia telangiectasia](#)
- [Atherosclerosis](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chromosome Aberrations](#)
- [Cognition](#)
- [Colon cancer](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Coronary Restenosis](#)
- [Diabetes Mellitus](#)

- [Disease Progression](#)
- [Edema](#)
- [Endometriosis](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Graft Occlusion](#)
- [Head and Neck Neoplasms](#)
- [Hyperparathyroidism](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Mouth Neoplasms](#)
- [Multiple endocrine neoplasia](#)
- [Multiple Endocrine Neoplasia Type 1](#)
- [Myocardial Infarction](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)

- [Parathyroid Neoplasms](#)
- [Precancerous Conditions](#)
- [Primary Ovarian Insufficiency](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)