


KLK3 monoclonal antibody, clone A67-B/E3 + 1A7

Catalog # MAB21091

Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Prostate Carcinoma using KLK3 monoclonal antibody, clone A67-B/E3 + 1A7.

Specification

Product Description Mouse monoclonal antibody raised against human KLK3.

Immunogen PSA (p30) from human sperm plasma

Host Mouse

Reactivity Human

Form Liquid

Purification Protein A/G purification

Isotype IgG

Recommend Usage
Flow Cytometry (0.5-1 ug/10⁶ cells in 0.1 mL)
Immunofluorescence (0.5-1ug/mL)
Immunohistochemistry (Formalin-fixed) (0.5-1 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer In 10 mM PBS (0.05% BSA, 0.05% sodium azide)

Storage Instruction Store at 2 to 8°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Prostate Carcinoma using KLK3 monoclonal antibody, clone A67-B/E3 + 1A7.

- Immunofluorescence
- Flow Cytometry

Gene Info — KLK3

Entrez GeneID [354](#)

Protein Accession# [P07288](#)

Gene Name KLK3

Gene Alias APS, KLK2A1, PSA, hK3

Gene Description kallikrein-related peptidase 3

Omim ID [176820](#)

Gene Ontology [Hyperlink](#)

Gene Summary Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Its protein product is a protease present in seminal plasma. It is thought to function normally in the liquefaction of seminal coagulum, presumably by hydrolysis of the high molecular mass seminal vesicle protein. Serum level of this protein, called PSA in the clinical setting, is useful in the diagnosis and monitoring of prostatic carcinoma. Alternate splicing of this gene generates several transcript variants encoding different isoforms. [provided by RefSeq]

Other Designations P-30 antigen|gamma-seminoprotein|kallikrein 3, (prostate specific antigen)|prostate specific antigen|semenogelase|seminin

Pathway

- [Pathways in cancer](#)
- [Prostate cancer](#)

Disease

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lymphatic Metastasis](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)