


IL7R monoclonal antibody, clone IL7R/2751

Catalog # MAB21102 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Tonsil using IL7R monoclonal antibody, clone IL7R/2751.

Specification

Product Description Mouse monoclonal antibody raised against partial recombinant human IL7R.

Immunogen Recombinant protein corresponding to amino acid 21-236 of human IL7R.

Host Mouse

Reactivity Human

Form Liquid

Purification Protein A/G purification

Isotype IgG2c, kappa

Recommend Usage Immunohistochemistry (Formalin-fixed) (2-4 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer In 10 mM PBS.

Storage Instruction Store at -20 to -80°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Tonsil using IL7R monoclonal antibody, clone IL7R/2751.

Gene Info — IL7R

Entrez GeneID [3575](#)

Protein Accession# [P16871](#)

Gene Name IL7R

Gene Alias CD127, CDW127, IL-7R-alpha, IL7RA, ILRA

Gene Description interleukin 7 receptor

Omim ID [146661 600802](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a receptor for interleukine 7 (IL7). The function of this receptor requires the interleukin 2 receptor, gamma chain (IL2RG), which is a common gamma chain shared by the receptors of various cytokines, including interleukine 2, 4, 7, 9, and 15. This protein has been shown to play a critical role in the V(D)J recombination during lymphocyte development. This protein is also found to control the accessibility of the TCR gamma locus by STAT5 and histone acetylation. Knockout studies in mice suggested that blocking apoptosis is an essential function of this protein during differentiation and activation of T lymphocytes. The functional defects in this protein may be associated with the pathogenesis of the severe combined immunodeficiency (SCID). [provided by RefSeq]

Other Designations CD127 antigen|interleukin 7 receptor alpha chain

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Hematopoietic cell lineage](#)
- [Jak-STAT signaling pathway](#)

- [Primary immunodeficiency](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bronchial Hyperreactivity](#)
- [Carcinoma](#)
- [Celiac Disease](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Graft vs Host Disease](#)
- [Helicobacter Infections](#)
- [HIV Infections](#)
- [Hyperparathyroidism](#)
- [Kaposi Varicelliform Eruption](#)
- [Lung Diseases](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphomatoid Granulomatosis](#)
- [Lymphopenia](#)
- [Multiple Sclerosis](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Hypersensitivity](#)
- [Sarcoidosis](#)

- [Severe combined immunodeficiency](#)
- [Stomach Neoplasms](#)
- [Syndrome](#)
- [Tobacco Use Disorder](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)