# KRT16 monoclonal antibody, clone R20-S

Catalog # MAB23284 Size 7 mL

## Applications



#### Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human skin sweat gland with KRT16 monoclonal antibody, clone R20-S (Cat # MAB23284).

Specification	
Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human KRT16.
Immunogen	A synthetic peptide corresponding to C-terminus of human KRT16.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	immunoaffinity purification
Recommend Usage	Immunohistochemistry (1:100-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In 20 mM Tris-HCI buffer, pH 8.0 (20 mg/mL BSA, 0.05% Sodium Azide).
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

😵 Abnova

### Applications

• Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human skin sweat gland with KRT16 monoclonal antibody, clone R20-S (Cat # MAB23284).

## Gene Info — KRT16

Entrez GenelD	3868
Protein Accession#	<u>P08779</u>
Gene Name	KRT16
Gene Alias	CK16, K16, K1CP, KRT16A, NEPPK
Gene Description	keratin 16
Omim ID	<u>144200 148067 167200 600962</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the keratin gene family. The keratins are interm ediate filament proteins responsible for the structural integrity of epithelial cells and are subdivide d into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins whic h are arranged in pairs of heterotypic keratin chains and are clustered in a region of chromosome 17q12-q21. This keratin has been coexpressed with keratin 14 in a number of epithelial tissues, i ncluding esophagus, tongue, and hair follicles. Mutations in this gene are associated with type 1 p achyonychia congenita, non-epidermolytic palmoplantar keratoderma and unilateral palmoplantar verrucous nevus. [provided by RefSeq
Other Designations	cytokeratin 16 focal non-epidermolytic palmoplantar keratoderma keratin, type I cytoskeletal 16