


HIP1 monoclonal antibody, clone 1B11

Catalog # MAB2341 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot Analysis of HIP1 in HeLa whole cell extracts using HIP1 monoclonal antibody, clone 1B11 (Cat # MAB2341).

Specification	
Product Description	Mouse monoclonal antibody raised against HIP1.
Immunogen	Human 3'-HIP1.
Host	Mouse
Theoretical MW (kDa)	65
Reactivity	Human, Mouse
Specificity	This antibody is specific to human HIP1. Does not cross-react with HIP1r.
Form	Liquid
lsotype	lgG1, kappa
Recommend Usage	Western Blot (1:10000) The optimal working dilution should be determined by the end user.
Storage Buffer	In buffer containing 0.09% sodium azide
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western Blot Analysis of HIP1 in HeLa whole cell extracts using HIP1 monoclonal antibody, clone 1B11 (Cat # MAB2341).

Gene Info — HIP1

Entrez GenelD	<u>3092</u>
Protein Accession#	<u>000291</u>
Gene Name	HIP1
Gene Alias	ILWEQ, MGC126506
Gene Description	huntingtin interacting protein 1
Omim ID	<u>176807 601767</u>
Gene Ontology	Hyperlink
Gene Summary	The product of this gene is a membrane-associated protein that colocalizes with huntingtin. This p rotein has similarities to cytoskeleton proteins and its interaction with huntingtin is thought to play a functional role in the cell filament network. Loss of normal huntingtin-HIP1 interaction in Huntingt on disease may contribute to a defect in membrane-cytoskeletal integrity in the brain. This gene c ould help in the understanding of the normal function of huntingtin and also the pathogenesis of Hu ntington disease. It also has been implicated in the pathogenesis of hematopoietic malignancies. An alternative splice variant of this gene has been described but its full length sequence has not b een determined. [provided by RefSeq
Other Designations	_

Other Designations

Publication Reference

Huntingtin interacting protein 1 modulates the transcriptional activity of nuclear hormone receptors.

Mills IG, Gaughan L, Robson C, Ross T, McCracken S, Kelly J, Neal DE.

The Journal of Cell Biology 2005 Jul; 170(2):191.

Application: ChIP, IF, IP, WB-Ce, WB-Tr, Human, Monkey, COS-7, LNCaP cells



Huntingtin-interacting protein 1 is overexpressed in prostate and colon cancer and is critical for cellular survival.

Rao DS, Hyun TS, Kumar PD, Mizukami IF, Rubin MA, Lucas PC, Sanda MG, Ross TS.

The Journal of Clinical Investigation 2002 Aug; 110(3):351.

Application: IHC-P, WB-Ce, Human, Mouse, Colon cancers, Human cancer cell lines, Human tissues, Prostate cancers

Huntingtin interacting protein 1 ls a clathrin coat binding protein required for differentiation of late spermatogenic progenitors.

Rao DS, Chang JC, Kumar PD, Mizukami I, Smithson GM, Bradley SV, Parlow AF, Ross TS.

Molecular and Cellular Biology 2001 Nov; 21(22):7796.

Application: IF, IHC, IP, WB-Ti, WB-Tr, Human, Mouse, Brains, Testes, HEK 293T cells, Seminiferous tubules

Disease

Huntington disease