

# KLK3 monoclonal antibody, clone 131-14234

Catalog # MAB4310

Size 1 mg

## Specification

**Product Description** Mouse monoclonal antibody raised against native KLK3.

**Immunogen** Native purified human KLK3.

**Host** Mouse

**Reactivity** Human

**Specificity** human Prostate Specific Antigen (PSA). PSA is a tumor-associated antigen with a molecular weight of 34 KDa. The antigen is found in normal prostatic epithelial lining cells of the ducts and acini, as well as in primary and metastatic prostate carcinoma cells. PSA reactivity may also be found in BPH (benign prostatic hypertrophy), though the pattern of staining is different than that found in carcinoma tissues. The staining pattern ranges from cytoplasmic and secretory in normal and benign prostate to cytoplasmic with membrane in neoplastic prostate.

**Form** Liquid

**Isotype** IgG1, kappa

**Recommend Usage** The optimal working dilution should be determined by the end user.

**Storage Buffer** In 10 mM phosphate buffer, pH 7.4 (150 mM NaCl, 0.09% sodium azide)

**Storage Instruction** Store at 4°C. For long term storage store at -20°C.  
Aliquot to avoid repeated freezing and thawing.

**Note** This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
- Immunohistochemistry (Frozen sections)

## Gene Info — KLK3

Entrez GeneID	<a href="#">354</a>
Gene Name	KLK3
Gene Alias	APS, KLK2A1, PSA, hK3
Gene Description	kallikrein-related peptidase 3
Omim ID	<a href="#">176820</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Its protein product is a protease present in seminal plasma. It is thought to function normally in the liquefaction of seminal coagulum, presumably by hydrolysis of the high molecular mass seminal vesicle protein. Serum level of this protein, called PSA in the clinical setting, is useful in the diagnosis and monitoring of prostatic carcinoma. Alternate splicing of this gene generates several transcript variants encoding different isoforms. [provided by RefSeq]
Other Designations	P-30 antigen gamma-seminoprotein kallikrein 3, (prostate specific antigen) prostate specific antigen semenogelase seminin

## Pathway

- [Pathways in cancer](#)
- [Prostate cancer](#)

## Disease

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)

- [Genetic Predisposition to Disease](#)
- [Lymphatic Metastasis](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)