IL2RA monoclonal antibody, clone MEM-181 (Biotin)

Catalog # MAB4577 Size 100 ug

Specification

Product Description	Mouse monoclonal antibody raised against native IL2RA.
Immunogen	Native purified IL2RA from PHA-activated peripheral blood leucocytes.
Host	Mouse
Theoretical MW (kDa)	55
Reactivity	Human
Specificity	This antibody reacts with CD25 (Interleukin-2 receptor alpha chain), a 55 KDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4+ T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes.
Form	Liquid
Conjugation	Biotin
Isotype	IgG1
Recommend Usage	Flow Cytometry (1:500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (0.09% sodium azide)
Storage Instruction	Store at 4°C. Do not freeze. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Info — IL2RA

Entrez GeneID	3559
Gene Name	IL2RA
Gene Alias	CD25, IDDM10, IL2R, TCGFR
Gene Description	interleukin 2 receptor, alpha
Omim ID	147730 601942
Gene Ontology	Hyperlink
Gene Summary	The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity r eceptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determine d to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated, b ut the significance of each is presently unknown. [provided by RefSeq
Other Designations	Interleukin-2 receptor|OTTHUMP00000019030|OTTHUMP00000044870|interleukin 2 receptor, al pha chain

Publication Reference

- CD25 indicates the neoplastic phenotype of mast cells: a novel immunohistochemical marker for the diagnosis of systemic mastocytosis (SM) in routinely processed bone marrow biopsy specimens.

  Sotlar K, Horny HP, Simonitsch I, Krokowski M, Aichberger KJ, Mayerhofer M, Printz D, Fritsch G, Valent P.

  The American Journal of Surgical Pathology 2004 Oct; 28(10):1319.

- Advances in interleukin 2 receptor targeted treatment.

  Morris JC, Waldmann TA.

  Annals of the Rheumatic Diseases 2000 Nov; 59:i109.

- The interleukin-2 receptor in human monocytes and macrophages: regulation of expression and release of the alpha and beta chains (p55 and p75).

  C Scheibenbogen, U Keilholz, M Richter, R Andreesen, W Hunstein.

  Research in Immunology 1992 Jan; 143(1):33.

  Application: Flow Cyt, Human, Human macrophages, Human monocytes

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Pathway

- Cytokine-cytokine receptor interaction
- Endocytosis
- Hematopoietic cell lineage
- Jak-STAT signaling pathway

Disease

- Acquired Immunodeficiency Syndrome
- Addison Disease
- Alopecia Areata
- Alzheimer Disease
- Arthritis
- Asthma
- Atrial Fibrillation
- Autoimmune Diseases
- Bronchiolitis
- Carcinoma
- Cardiovascular Diseases
- Celiac Disease
- Chorioamnionitis
- Colitis
- Connective Tissue Diseases
- Crohn Disease
- Diabetes Mellitus
- Disease Progression

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Edema
- Fetal Diseases
- Fetal Membranes
- Genetic Predisposition to Disease
- Graves Disease
- Hyperparathyroidism
- Hypersensitivity
- Infant
- Infection
- Inflammation
- Lupus Erythematosus
- Measles
- Melanoma
- Multiple Sclerosis
- Mumps
- Musculoskeletal Diseases
- Obstetric Labor
- Prediabetic State
- Pre-Eclampsia
- Pregnancy Complications
- Premature Birth
- Prostatic Neoplasms
- Respiratory Syncytial Virus Infections
- Rubella
- Skin Diseases
- Thyroid Diseases

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Vasculitis
- Vitiligo

Copyright © 2023 Abnova Corporation. All Rights Reserved.