


# Tnfrsf8 monoclonal antibody, clone 2SH12-5F (PE)

Catalog # MAB5711 Size 100 ug

## **Applications**



#### Flow Cytometry

The YAC-1 cell line was stained with Tnfrsf8 monoclonal antibody, clone 2SH12-5F (FITC) (Cat # MAB5709). Large cells were gated and analyzed by flow cytometry.

Specification	
Product Description	Hamster monoclonal antibody raised against partial recombinant Tnfrsf8.
Immunogen	Recombinant mouse IgG1 fusion protein corresponding to extracellular region of mouse Tnfrsf8.
Host	Hamster
Reactivity	Mouse
Specificity	Murine CD30, Mr 52 KDa (core peptide).
Form	Liquid
Conjugation	PE
Isotype	lgG, kappa
Recommend Usage	Flow Cytometry (0.2 ug/10 <sup>6</sup> cells)  The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)



#### **Product Information**

Storage Instruction	Store in the dark at 4°C. Do not freeze.  Avoid prolonged exposure to light.  Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## **Applications**

Flow Cytometry

The YAC-1 cell line was stained with Tnfrsf8 monoclonal antibody, clone 2SH12-5F (FITC) (Cat # MAB5709). Large cells were gated and analyzed by flow cytometry.

Gene Info — Tnfrsf8	
Entrez GeneID	<u>21941</u>
Gene Name	Tnfrsf8
Gene Alias	Cd30, D1S166E, Ki, Ki-1
Gene Description	tumor necrosis factor receptor superfamily, member 8
Gene Ontology	<u>Hyperlink</u>
Gene Summary	0
Other Designations	OTTMUSP00000010648

#### **Publication Reference**

• Structure and expression of murine CD30 and its role in cytokine production.

Bowen MA, Lee RK, Miragliotta G, Nam SY, Podack ER.

The Journal of Immunology 1996 Jan; 156(2):442.

• Regulation of murine B cell growth and differentiation by CD30 ligand.

Shanebeck KD, Maliszewski CR, Kennedy MK, Picha KS, Smith CA, Goodwin RG, Grabstein KH.

European Journal of Immunology 1995 Aug; 25(8):2147.

Application: Flow Cyt, Func, Mouse, B cells, KO3 cells, T cells



## **Product Information**

• CD30 (Ki-1) molecule: a new cytokine receptor of the tumor necrosis factor receptor superfamily as a tool for diagnosis and immunotherapy.

Falini B, Pileri S, Pizzolo G, Durkop H, Flenghi L, Stirpe F, Martelli MF, Stein H. Blood 1995 Jan; 85(1):1.