
CD81 monoclonal antibody, clone M38 (FITC)

Catalog # MAB6494 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric surface staining of human peripheral blood stained with CD81 monoclonal antibody, clone M38 (FITC) (Cat # MAB6494).

Specification

Product Description	Mouse monoclonal antibody raised against native CD81.
Immunogen	Native purified CD81 from human T-ALL cell line (MOLT-4).
Host	Mouse
Theoretical MW (kDa)	25
Reactivity	Cat, Human, Rabbit
Specificity	This antibody reacts with CD81, a 25 KDa member of the tetraspanin family, expressedon majority o f cells.
Form	Liquid
Conjugation	FITC
lsotype	lgG1
Recommend Usage	The optimal working dilution should be determined by the end user.

😵 Abnova

Product Information

Storage Buffer	In PBS (0.2% BSA, 0.09% sodium azide)
Storage Instruction	Store in the dark at 4°C. Do not freeze. Avoid prolonged exposure to light. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

• Flow Cytometry

Flow cytometric surface staining of human peripheral blood stained with CD81 monoclonal antibody, clone M38 (FITC) (Cat # MAB6494).

Gene Info — CD81

Entrez GenelD	<u>975</u>
Gene Name	CD81
Gene Alias	S5.7, TAPA1, TSPAN28
Gene Description	CD81 molecule
Omim ID	<u>186845</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known a s the tetraspanin family. Most of these members are cell-surface proteins that are characterized b y the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded prot ein is a cell surface glycoprotein that is known to complex with integrins. This protein appears to p romote muscle cell fusion and support myotube maintenance. Also it may be involved in signal transduction. This gene is localized in the tumor-suppressor gene region and thus it is a candidate g ene for malignancies. [provided by RefSeq
Other Designations	26 kDa cell surface protein TAPA-1 CD81 antigen CD81 antigen (target of antiproliferative antibo dy 1) target of antiproliferative antibody 1

Publication Reference

😵 Abnova

Product Information

 <u>Selective enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and on</u> <u>exosomes secreted by human B-lymphocytes.</u>

Escola JM, Kleijmeer MJ, Stoorvogel W, Griffith JM, Yoshie O, Geuze HJ. The Journal of Biological Chemistry 1998 Aug; 273(32):20121.

 Molecular analyses of the association of CD4 with two members of the transmembrane 4 superfamily, CD81 and CD82.

T Imai, M Kakizaki, M Nishimura, O Yoshie. Journal of Immunology 1995 Aug; 155(3):1229.

 C33 antigen and M38 antigen recognized by monoclonal antibodies inhibitory to syncytium formation by human T cell leukemia virus type 1 are both members of the transmembrane 4 superfamily and associate with each other and with CD4 or CD8 in T cells.

Imai T, Yoshie O.

The Journal of Immunology 1993 Dec; 151(11):6470.

Pathway

<u>B cell receptor signaling pathway</u>

Disease

- Atherosclerosis
- <u>Carcinoma</u>
- Genetic Predisposition to Disease
- Hematologic Diseases
- Hepatitis C
- Hodgkin Disease
- Kidney Failure
- Liver Neoplasms
- Lung Neoplasms
- Lymphoproliferative Disorders

😵 Abnova

- <u>Multiple Myeloma</u>
- <u>Obesity</u>
- <u>Occupational Diseases</u>
- Pulmonary Disease
- <u>Urinary Bladder Neoplasms</u>
- <u>Waldenstrom Macroglobulinemia</u>
- Werner syndrome