
KRT18 monoclonal antibody, clone DA-7 (Biotin)

Catalog # MAB6506 Size 100 ug

Applications



Western Blot (Cell lysate)

Western Blotting analysis (reducing conditions) of whole cell lysate using KRT18 monoclonal antibody, clone DA-7 (Cat # MAB0860).
Lane 1 : HeLa (human cervix carcinoma cell line).
Lane 2 : A-431 (human epidermoid carcinoma cell line).
Lane 3 : MCF-7 (human breast adenocarcinoma cell line).

Specification

Product Description	Mouse monoclonal antibody raised against native KRT18.
Immunogen	Native purified KRT18 from human breast carcinoma cell line (PMC-42).
Host	Mouse
Theoretical MW (kDa)	45
Reactivity	Human
Specificity	This antibody reacts with Cytokeratin 18 (45 KDa).
Form	Liquid
Conjugation	Biotin
Isotype	IgG1
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.09% sodium azide)

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at 4°C. Do not freeze. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
- Immunocytochemistry
- Enzyme-linked Immunoabsorbent Assay
- Immunoprecipitation
- Western Blot (Cell lysate)

  Western Blotting analysis (reducing conditions) of whole cell lysate using KRT18 monoclonal antibody, clone DA-7 (Cat # MAB0860).
  Lane 1 : HeLa (human cervix carcinoma cell line).
  Lane 2 : A-431 (human epidermoid carcinoma cell line).
  Lane 3 : MCF-7 (human breast adenocarcinoma cell line).

Gene Info — KRT18

Entrez GeneID	3875
Gene Name	KRT18
Gene Alias	CYK18, K18
Gene Description	keratin 18
Omim ID	148070 215600
Gene Ontology	Hyperlink
Gene Summary	KRT18 encodes the type I intermediate filament chain keratin 18. Keratin 18, together with its fila ment partner keratin 8, are perhaps the most commonly found members of the intermediate filam ent gene family. They are expressed in single layer epithelial tissues of the body. Mutations in this gene have been linked to cryptogenic cirrhosis. Two transcript variants encoding the same protei n have been found for this gene. [provided by RefSeq
Other Designations	cell proliferation-inducing protein 46|cytokeratin 18

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Pathway

- Pathogenic Escherichia coli infection - EHEC

Disease

- Cleft Lip
- Cleft Palate
- Drug-Induced Liver Injury
- Genetic Predisposition to Disease
- Liver Cirrhosis
- Liver Failure

Copyright © 2023 Abnova Corporation. All Rights Reserved.