

Bioactive

PDGFB (Human) Recombinant Protein

Catalog # P6219

Size 100 ug

Applications

Result of activity analysis

Result of activity analysis

□

Specification

Product Description	Human PDGFB (P01127) recombinant protein expressed in <i>E.Coli</i> .
Sequence	MSLGSLTIAEPAMAECKTRTEVFEISRRLIDRTNANFLVWPPCVEVQRCSGCCNNRNVQCRPTQ VQLRPVQVRKIEVRKKPIFKKATVTLEDHLACKCETVAAARPVT
Host	Escherichia coli
Theoretical MW (kDa)	12.4/24.9
Form	Lyophilized
Purity	>= 95%
Endotoxin Level	<= 1 EUs/ug (Kinetic LAL)
Activity	ED ₅₀ <= 20 ng/mL NR6R-3T3 cell proliferation The values provided above are minimum expected values to pass internal requirements.
Quality Control Testing	Reducing and Non-Reducing SDS PAGE
Conformation	Dimer

Storage Buffer	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate , pH 7.5.
Storage Instruction	Stored at -20°C to -80°C for 12 month. After reconstitution with sterile water at 0.1 mg/mL, store at -20°C to -80°C for 3 months, store at 4°C for 1 month. Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis Result of activity analysis

Applications

- Western Blot
- Functional Study

Gene Info — PDGFB

Entrez GeneID	5155
Protein Accession#	P01127
Gene Name	PDGFB
Gene Alias	FLJ12858, PDGF2, SIS, SSV, c-sis
Gene Description	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)
Omim ID	190040
Gene Ontology	Hyperlink

Gene Summary	The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor. Two alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq]
--------------	--

Other Designations

PDGF, B chain|Platelet-derived growth factor, beta polypeptide (oncogene SIS)|becaplermin|oncogene SIS|platelet-derived growth factor 2|platelet-derived growth factor beta|platelet-derived growth factor, B chain|v-sis platelet-derived growth factor beta p

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Focal adhesion](#)
- [Gap junction](#)
- [Glioma](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Renal cell carcinoma](#)

Disease

- [Amyotrophic lateral sclerosis](#)
- [Anoxia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Fetal Membranes](#)

- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Head and Neck Neoplasms](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Ovarian Neoplasms](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Recurrence](#)