Bioactive

SHH (Human) Recombinant Protein

Catalog # P6402 Size 25 ug

Specification

Product Description	Human SHH (Q15465) recombinant protein expressed in *E. Coli*.
Sequence	IVIGPGRGFGKRRHPKKLTPLAYKQFIPNVAEKTLGASGRYEGKISRNSERFKELTPNYNPDIIFKDE ENTGADRLMTQRCKDKLNALAISVMNQWPGVKLRVTEGWDEDGHHSEESLHYEGRALDITTSDR DRSKYGMLARLAVEAGFDWVYYESKAHIHCSVKAENSVAAKSGG
Host	Escherichia coli
Theoretical MW (kDa)	19
Form	Lyophilized
Purity	> 95%
Endotoxin Level	<= 1 EUs/ug (LAL gel clot method)
Activity	Determined by the ability to induce alkaline phosphatase production by C3H/10T1/2 (CCL-226) cells . The expected ED_{50} for this effect is 0.8-1 ug/mL.
Storage Buffer	Lyophilized from PBS, pH 7.2.
Storage Instruction	Stored at -20°C to-80°C. After reconstitution with sterile water not less than 0.1 mg/mL, store at -20°C to -80°C for 6 months, st ore at 4°C for 1 month. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- Western Blot

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Info — SHH

Entrez GeneID	6469
Protein Accession#	Q15465
Gene Name	SHH
Gene Alias	HHG1, HLP3, HPE3, MCOPCB5, SMMCI, TPT, TPTPS
Gene Description	sonic hedgehog homolog (Drosophila)
Omim ID	120200 142945 147250 174500 600725
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a protein that is instrumental in patterning the early embryo. It has been implicated as the key inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Of three human proteins showing sequence and functional similarity to the sonic hedgehog protein of Drosophila, this protein is the most similar. The protein is made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the developing embryo. Defects in this protein or in its signalling pathway are a cause of holoprosencephaly (HPE), a disorder in which the developing forebrain fails to correctly separate into right and left hemispheres. HPE is manifested by facial deformities. It is also thought that mutations in this gene or in its signalling pathway may be responsible for VACTERL syndrome, which is characterized by vertebral defects, anal atresia, tracheoesophageal fistula with esophageal atresia, radial and renal dysplasia, cardiac anomalies, and limb abnormalities. Additionally, mutations in a long range enhancer located approximately 1 megabase upstream of this gene disrupt limb patterning and can result in preaxial polydactyly. [provided by RefSeq
Other Designations	sonic hedgehog

Pathway

- Basal cell carcinoma
- Hedgehog signaling pathway
- Pathways in cancer

Disease

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Cleft Lip
- Cleft Palate
- Genetic Predisposition to Disease
- Holoprosencephaly
- Kidney Failure
- Parkinson disease
- Sleep Apnea
- Syndrome
- Thyroid Neoplasms

Copyright © 2023 Abnova Corporation. All Rights Reserved.