


Bioactive

HuPro®

TNFRSF17 (Human) Recombinant Protein

Catalog # P6735 Size 100 ug

Applications



Result of activity analysis

Result of activity analysis

Specification	
Product Description	Human TNFRSF17 (NP_001183.2, 1 a.a 54 a.a.) partial recombinant protein with a Fc, 6×His tag at the C-terminus expressed in HEK293 cells.
Host	Human
Form	Lyophilized
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Ni-sepharose purification
Purity	> 92% (determined by SDS-PAGE)
Endotoxin Level	< 0.1 EU/ug of protein (determined by LAL method)



Product Information

Activity	Measured by the binding ability in a functional ELISA. Immobilized recombinant human TNFSF13B a t 5 ug/mL (100 uL/well) can bind recombinant human TNFRSF17 with a linear range of 3-20 ng/mL.
Quality Control Testing	SDS-PAGE Stained with Coomassie Blue
Recommend Usage	SDS-PAGE
	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from a 0.22 um filtered solution of PBS, pH 7.4.
Storage Instruction	Store at -80°C on dry atmosphere, lyophilized antibodies are stable at least 2 years.
	After reconstitution with deionized water, store at -20°C or lower.
	Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis
	Result of activity analysis

Applications

SDS-PAGE

Gene Info — TNFRSF17	
Entrez GenelD	<u>608</u>
Protein Accession#	Q02223
Gene Name	TNFRSF17
Gene Alias	BCM, BCMA, CD269
Gene Description	tumor necrosis factor receptor superfamily, member 17
Omim ID	<u>109545</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development a nd autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq
Other Designations	B cell maturation antigen B-cell maturation factor OTTHUMP00000160261



Pathway

Cytokine-cytokine receptor interaction

Disease

- Arthritis
- Asthma
- Colitis
- Crohn Disease
- Genetic Predisposition to Disease
- Hematologic Diseases
- Kidney Failure
- Multiple Myeloma
- Occupational Diseases