

Bioactive


HuPro®

TNFRSF8 (Human) Recombinant Protein

Catalog # P6749

Size 100 ug

Applications



Result of activity analysis

Result of activity analysis

Specification

Product Description	Human TNFRSF8 (NP_001234.2, 19 a.a. - 379 a.a.) partial recombinant protein with a 6xHis tag at the C-terminus expressed in HEK293 cells.
Host	Human
Form	Lyophilized
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Ni-sepharose purification
Purity	> 97% (determined by SDS-PAGE)
Endotoxin Level	< 0.1 EU/ug of protein (determined by LAL method)

Activity	Measured by the binding ability in a functional ELISA. Immobilized recombinant human TNFRSF8 at 2 ug/mL (100 uL/well) can bind recombinant human TNFSF8 with a linear range of 1.22-15.23 ng/mL.
Quality Control Testing	SDS-PAGE Stained with Coomassie Blue
Recommend Usage	SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from a 0.22 um filtered solution of PBS, pH 7.4.
Storage Instruction	Store at -80°C on dry atmosphere, lyophilized antibodies are stable at least 2 years. After reconstitution with deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis Result of activity analysis

Applications

- SDS-PAGE

Gene Info — TNFRSF8

Entrez GeneID	943
Protein Accession#	P28908
Gene Name	TNFRSF8
Gene Alias	CD30, D1S166E, KI-1
Gene Description	tumor necrosis factor receptor superfamily, member 8
Omim ID	153243
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]

Other Designations

CD30 antigen|CD30L receptor|Ki-1 antigen|OTTHUMP00000001783|cytokine receptor CD30|lymphocyte activation antigen CD30

Pathway

- [Cytokine-cytokine receptor interaction](#)

Disease

- [Asthma](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [HIV Infections](#)
- [Kidney Failure](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Tobacco Use Disorder](#)