Bioactive

PDFGB (Human) Recombinant Protein

Catalog # P7140 Size 10 ug

Applications



Result of bioactivity analysis

Result of bioactivity analysis

Specification

Product Description	Human PDGFB (P01127, 82 a.a. - 190 a.a.) partial recombinant protein expressed in *Pichia pastoris*.
Sequence	SLGSLTIAEPAMIAECKTRTEVFEISRRLIDRTNANFLVWPPCVEVQRCSGCCNNRNVQCRPTQV QLRPVQVRKIEIVRKKPIFKKATVTLEDHLACKCETVAAARPVTRSPGGSQEQR
Host	Yeast
Theoretical MW (kDa)	24.3
Form	Lyophilized
Preparation Method	*Pichia pastoris* expression system
Purity	> 97% by SDS-PAGE
Endotoxin Level	< 1 EU per 1 ug of protein (determined by gel clotting method)
Activity	The ED_{50} was determined by the dose-dependent stimulation of proliferation of Balb/c 3T3 cells is 5.0 - 10.0 ng/ml, corresponding to a specific activity of > 3.3 x 10^5 units/mg.
Storage Buffer	Lyophilized from sterile distilled Water up to 100 ug/ml

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at 4°C to 8°C for 1 week. For long term storage store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Result of bioactivity analysis Result of bioactivity analysis

Applications

- Functional Study
- SDS-PAGE

Gene Info — PDGFB

Entrez GeneID	5155
Protein Accession#	P01127
Gene Name	PDGFB
Gene Alias	FLJ12858, PDGF2, SIS, SSV, c-sis
Gene Description	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)
Omim ID	190040
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the platelet-derived growth factor family. The fou r members of this family are mitogenic factors for cells of mesenchymal origin and are characteriz ed by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) o r as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where th e dimers are connected by disulfide bonds. Mutations in this gene are associated with meningio ma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosa rcoma protuberans resulting from unregulated expression of growth factor. Two alternatively splice d transcript variants encoding different isoforms have been identified for this gene. [provided by R efSeq
Other Designations	PDGF, B chain|Platelet-derived growth factor, beta polypeptide (oncogene SIS)|becaplermin|onc ogene SIS|platelet-derived growth factor 2|platelet-derived growth factor beta|platelet-derived gro wth factor, B chain|v-sis platelet-derived growth factor beta p

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Cytokine-cytokine receptor interaction
- Focal adhesion
- Gap junction
- Glioma
- MAPK signaling pathway
- Melanoma
- Pathways in cancer
- Prostate cancer
- Regulation of actin cytoskeleton
- Renal cell carcinoma

Disease

- Amyotrophic lateral sclerosis
- Anoxia
- Carcinoma
- Cardiovascular Diseases
- Chorioamnionitis
- Coronary Artery Disease
- Diabetes Mellitus
- Disease Progression
- Edema
- Fetal Membranes
- Genetic Predisposition to Disease
- Glomerulonephritis
- Head and Neck Neoplasms

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Neoplasms
- Obstetric Labor
- Ovarian Neoplasms
- Pre-Eclampsia
- Premature Birth
- Recurrence

Copyright © 2023 Abnova Corporation. All Rights Reserved.