

HuPro®

FGFR4 (Human) Recombinant Protein

Catalog # P7607

Size 50 ug

Specification

Product Description	Human FGFR4 (P22455, 22 a.a. - 369 a.a.) partial recombinant protein with hFc tag at C-terminus expressed in HEK293 cells.
Sequence	LEAEEVELEPCLAPSLEQQEQELTVALGQPVRLCCGRAERGGHWYKEGSRLAPAGRVRGWR GRLEIASFLPEDAGRYLCLARGSMVLQNLTLITGDSLTSNDDDEPKSHRDP SNRHSYPQQAPY WTHPQRMEKKLHAVPAGNTVKFRCPAAGNPTPTIRWLKDGQAFHGENRIGGIRLRHQHWSLVME SVVPSDRGTYTCLVENAVGSIRYNLLDVLERSPHRPILQAGLPANTTAVVGSDVELLCKVYSDAQ PHIQWLKHIVINGSSFGADGFPYVQVLKTADINSSEVEVLYLRNVSAEDAGEYTCLAGNSIGLSYQS AWLTVLPEEDPTWTAAAPARYTD
Host	Mammals
Theoretical MW (kDa)	76.3
Form	Lyophilized
Preparation Method	Mammalian cell (HEK293) expression system
Purity	> 90% by SDS-PAGE
Endotoxin Level	< 1 EU per 1 ug of protein (determined by gel clotting method)
Storage Buffer	Lyophilized from sterile distilled Water up to 100 ug/mL
Storage Instruction	Store at 4°C to 8°C for 1 week. For long term storage store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- SDS-PAGE

Gene Info — FGFR4

Entrez GeneID [2264](#)

Protein Accession# [P22455](#)

Gene Name FGFR4

Gene Alias CD334, JTK2, MGC20292, TKF

Gene Description fibroblast growth factor receptor 4

Omim ID [134935](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. The genomic organization of this gene, compared to members 1-3, encompasses 18 exons rather than 19 or 20. Although alternative splicing has been observed, there is no evidence that the C-terminal half of the IgIII domain of this protein varies between three alternate forms, as indicated for members 1-3. This particular family member preferentially binds acidic fibroblast growth factor and, although its specific function is unknown, it is overexpressed in gynecological tumor samples, suggesting a role in breast and ovarian tumorigenesis. [provided by RefSeq]

Other Designations OTTHUMP00000161430|hydroxyaryl-protein kinase|protein-tyrosine kinase|tyrosine kinase related to fibroblast growth factor receptor|tyrosylprotein kinase

Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [ACTH-Secreting Pituitary Adenoma](#)

- [Adenocarcinoma](#)
- [Bone Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Carcinoma](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colon cancer](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hypersensitivity](#)
- [Irritable Bowel Syndrome](#)
- [Kidney Failure](#)
- [Liver Neoplasms](#)
- [Lung carcinoma](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Mouth Neoplasms](#)
- [Neoplasm Metastasis](#)
- [Pituitary ACTH Hypersecretion](#)

- [Pituitary Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)
- [Recurrence](#)
- [Sarcoma](#)
- [Skin Neoplasms](#)
- [Soft Tissue Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Urinary Bladder Neoplasms](#)