


Bioactive

FGF1 (Human) Recombinant Protein

Catalog # P8115 Size 1 mg

Applications



SDS-PAGE analysis of FGF1 (Human) Recombinant Protein.

Specification

Product Description	Human FGF1 recombinant protein with polyhistidine tag at the C-terminus expressed in <i>Escherichia coli</i> .
Sequence	MFNLPPGNYKKPKLLYCSNGGHFLRLPDGTVDGTRDRSDQHIQLQLSAESVGEVYIKSTETGQYL AMDTDGLLYGSQTPNEECLFLERLEENHYNTYISKKHAEKNWVGLKKNKGSCCKRGPRTHYGQKAI LFLPLPVSSD with polyhistidine tag at the C-terminus.
Host	Escherichia coli
Form	Lyophilized
Preparation Method	<i>Escherichia coli</i> expression system
Purity	> 98% as determined by SDS-PAGE.
Endotoxin Level	< 0.1 EU per 1 ug of the protein by the LAL method.
Activity	Measured by the induction of 3T3 cells proliferation. The ED ₅₀ for this effect is < 0.3 ng/mL. The specific activity of recombinant human FGF-1 is > 1 x 10 ⁶ IU/mg.
Quality Control Testing	SDS-PAGE Stained with Coomassie Blue. SDS-PAGE analysis of FGF1 (Human) Recombinant Protein.

Recommend Usage	SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from a solution containing 1X PBS, pH 8.0. Reconstitute the lyophilized powder in ddH ₂ O to 100 ug/mL.
Storage Instruction	Lyophilized protein should be stored at -20°C. Protein aliquots should be stored at -20°C to -80°C. This product is stable for one year. Avoid repeated freeze/thaw cycles.

Applications

- Functional Study
- SDS-PAGE

Gene Info — FGF1

Entrez GeneID	2246
Gene Name	FGF1
Gene Alias	AFGF, ECGF, ECGF-beta, ECGFA, ECGFB, FGF-alpha, FGFA, GLIO703, HBGF1
Gene Description	fibroblast growth factor 1 (acidic)
Omim ID	131220
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Multiple alternatively spliced variants encoding different isoforms have been described. [provided by RefSeq]
Other Designations	OTTHUMP00000066028 OTTHUMP00000066030 OTTHUMP00000066031 OTTHUMP00000174675 endothelial cell growth factor, alpha endothelial cell growth factor, beta heparin-binding growth factor 1

Pathway

- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Alzheimer disease](#)
- [Arthritis](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Chorioamnionitis](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Diabetes Complications](#)
- [Endometriosis](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Hyperparathyroidism](#)
- [Hypertension](#)
- [Intracranial Aneurysm](#)
- [Metabolic Syndrome X](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Neovascularization](#)

- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Schizophrenia](#)