

Bioactive

CSF2 (Human) Recombinant Protein

Catalog # P8741 Size 2 x 10 ug

Specification

Product Description	Human CSF2 recombinant protein expressed in <i>Pichia Pastoris</i> .
Host	Yeast
Theoretical MW (kDa)	26-32
Form	Lyophilized
Preparation Method	yeast expression system
Purification	chromatographic
Purity	> 97% as determined by (a) RP-HPLC.(b) SDS-PAGE.
Activity	ED ₅₀ < 0.183 ng/mL, measured by the dose-dependant stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line), corresponding to a specific activity of 5.5 x 10 ⁶ IU/mg.
Storage Buffer	Protein (1 mg/mL) was lyophilized from a solution containing 10 mM phosphate buffer, pH 7.0, 40 mg mannitol, 10 mg sucrose. Reconstitute the lyophilized powder in ddH ₂ O to 100 ug/mL.
Storage Instruction	Lyophilized protein at room temperature for 3 weeks, should be stored at -20°C. Protein aliquots at 4 °C for 2-7 days and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid repeated freeze/thaw cycles.

Applications

- Functional Study

Gene Info — CSF2

Entrez GeneID	1437
Protein Accession#	P04141
Gene Name	CSF2
Gene Alias	GMCSF, MGC131935, MGC138897
Gene Description	colony stimulating factor 2 (granulocyte-macrophage)
Omim ID	138960
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q 31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. [provided by RefSeq]
Other Designations	colony stimulating factor 2 granulocyte-macrophage colony stimulating factor molgramostin sargramostim

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Fc epsilon RI signaling pathway](#)
- [Hematopoietic cell lineage](#)
- [Jak-STAT signaling pathway](#)
- [Natural killer cell mediated cytotoxicity](#)
- [T cell receptor signaling pathway](#)

Disease

- [Alzheimer disease](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Blindness](#)

- [Brain Ischemia](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chlamydia Infections](#)
- [Chorioamnionitis](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Dermatitis](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Erythema Nodosum](#)
- [Esophagitis](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Hypertension](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Metabolic Syndrome X](#)
- [Migraine Disorders](#)

- [Neoplasms](#)
- [Neovascularization](#)
- [Obesity](#)
- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Pain](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Pulmonary Disease](#)
- [Radiation Injuries](#)
- [Radiation Pneumonitis](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Rhinitis](#)
- [Sarcoidosis](#)
- [Stroke](#)
- [Thyroid Neoplasms](#)
- [Trachoma](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)