Bioactive

TNFSF10 (Human) Recombinant Protein

Catalog # P9294 Size 2 x 10 ug

Specification

Product Description	Human TNFSF10 partial recombinant protein expressed in?*Escherichia coli*.
Sequence	MVRERGPQRVAAHITGTRGRSNTLSSPNSKNEKALGRKINSWESSRSGHSFLSNLHLRNGELVIH EKGFYYIYSQTYFRFQEEIKENTKNDKQMVQYIYKYTSYPDPILLMKSARNSCWSKDAEYGLYSIYQG GIFELKENDRIFVSVTNEHLIDMDHEASFFGAFLVG
Host	Escherichia coli
Theoretical MW (kDa)	19.6
Form	Liquid
Preparation Method	*Escherichia coli* expression system
Purification	chromatographic
Purity	> 95% as determined by SDS-PAGE.
Activity	ED_{50} ≤ 1 ng/mL, measured in a cell cytotoxicity assay using Jurkat human T lymphocyte.
Storage Buffer	Solution (1 mg/mL) containing 20 mM Tris-HCl, pH 7.5, 10% glycerol, 300 mM NaCl, 0.1 mM DTT.
Storage Instruction	Store at 4°C for one weeks and should be stored at -20°C to -80°C. For long term storage it is reco mmended to add a carrier protein (0.1% HSA or BSA). Avoid repeated freeze/thaw cycles.

Applications

- Functional Study

Gene Info — TNFSF10

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Entrez GeneID	8743
Protein Accession#	P50591
Gene Name	TNFSF10
Gene Alias	APO2L, Apo-2L, CD253, TL2, TRAIL
Gene Description	tumor necrosis factor (ligand) superfamily, member 10
Omim ID	603598
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) lig and family. This protein preferentially induces apoptosis in transformed and tumor cells, but does not appear to kill normal cells although it is expressed at a significant level in most normal tissues. This protein binds to several members of TNF receptor superfamily including TNFRSF10A/TRAIL R1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and possibly also to TNFRSF11B/OPG. The activity of this protein may be modulated by binding to the decoy receptor s TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and TNFRSF11B/OPG that cannot induce apo ptosis. The binding of this protein to its receptors has been shown to trigger the activation of MAP K8/JNK, caspase 8, and caspase 3. [provided by RefSeq
Other Designations	Apo-2 ligand|TNF-related apoptosis inducing ligand TRAIL|tumor necrosis factor (ligand) family, member 10|tumor necrosis factor apoptosis-inducing ligand splice variant delta

Pathway

- Apoptosis
- Cytokine-cytokine receptor interaction
- Natural killer cell mediated cytotoxicity

Disease

- Breast Neoplasms
- Carcinoma
- Genetic Predisposition to Disease
- Head and Neck Neoplasms
- Hematologic Diseases

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Hodgkin Disease
- Kidney Failure
- Lupus Erythematosus
- Lymphoproliferative Disorders
- Multiple Myeloma
- Multiple Sclerosis
- Neoplasm Recurrence
- Neoplasms
- Occupational Diseases
- Ovarian Neoplasms
- Waldenstrom Macroglobulinemia
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.