


Bioactive

HuPro®

# FGFR2 (Human) Recombinant Protein


Catalog # P9927      Size 100 ug

## Applications



### Enzyme-linked Immunoabsorbent Assay

Immobilized Human FGFR2 alpha (IIIb), His Tag at 0.5 $\mu$ g/ml (100 $\mu$ l/well) on the plate. Dose response curve for Anti-FGFR2 alpha (IIIb) Antibody, hFc Tag with the EC50 of 6.1ng/ml determined by ELISA (QC Test).



### SEC-HPLC

The purity of Human FGFR2 alpha (IIIb) is greater than 95% as determined by SEC-HPLC.



### Tris-Bis PAGE

Human FGFR2 alpha (IIIb) on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.

## Specification

### Product Description

Human FGFR2 (P21802-3, Arg22-Glu378) partial recombinant protein with His tag at C-Terminus expressed in HEK293 cells.

<b>Sequence</b>	Arg22-Glu378
<b>Host</b>	Human
<b>Theoretical MW (kDa)</b>	40.7
<b>Form</b>	Lyophilized
<b>Preparation Method</b>	Mammalian cell (HEK293) expression system
<b>Purity</b>	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
<b>Endotoxin Level</b>	< 1 EU per 1 ug of protein (determined by LAL method)
<b>Activity</b>	The EC <sub>50</sub> was 6.1 ng/mL, measured by ELISA at 0.5 ug/mL.
<b>Quality Control Testing</b>	SEC-HPLC and Tris-Bis PAGE SEC-HPLC The purity of Human FGFR2 alpha (IIIb) is greater than 95% as determined by SEC-HPLC. Tris-Bis PAGE Human FGFR2 alpha (IIIb) on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%. %
<b>Recommend Usage</b>	Biological Activity ELISA SDS-PAGE The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	Lyophilized from sterile distilled Water is > 100 ug/mL
<b>Storage Instruction</b>	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	Result of bioactivity analysis

## Applications

- Enzyme-linked Immunoabsorbent Assay

Immobilized Human FGFR2 alpha (IIIb), His Tag at 0.5ug/ml (100ul/well) on the plate. Dose response curve for Anti-FGFR2 alpha (IIIb) Antibody, hFc Tag with the EC50 of 6.1ng/ml determined by ELISA (QC Test).

- Functional Study
- SDS-PAGE

## Gene Info — FGFR2

Entrez GeneID	<a href="#">2263</a>
Protein Accession#	<a href="#">P21802-3</a>
Gene Name	FGFR2
Gene Alias	BEK, BFR-1, CD332, CEK3, CFD1, ECT1, FLJ98662, JWS, K-SAM, KGFR, TK14, TK25
Gene Description	fibroblast growth factor receptor 2
Omim ID	<a href="#">101200</a> <a href="#">101400</a> <a href="#">101600</a> <a href="#">123150</a> <a href="#">123500</a> <a href="#">123790</a> <a href="#">137215</a> <a href="#">149730</a> <a href="#">176943</a> <a href="#">207410</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq]
Other Designations	BEK fibroblast growth factor receptor FGF receptor OTTHUMP00000020621 OTTHUMP00000020629 bacteria-expressed kinase hydroxyaryl-protein kinase keratinocyte growth factor receptor protein tyrosine kinase, receptor like 14 soluble FGFR4 variant 4

## Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)

## Disease

- [Acrocephalosyndactylia](#)
- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Craniofacial Dysostosis](#)
- [Craniosynostoses](#)
- [Cystadenocarcinoma](#)
- [Depressive Disorder](#)
- [Diabetes Complications](#)
- [Disease Susceptibility](#)
- [Endometrial Neoplasms](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Hypospadias](#)
- [Lymphatic Metastasis](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Metabolic Syndrome X](#)

- [Neoplasm Invasiveness](#)
- [Neoplasms](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Schizophrenia](#)
- [Skin Neoplasms](#)
- [Sleep Apnea](#)
- [Tobacco Use Disorder](#)