


CASP3 polyclonal antibody

Catalog # PAB0243


Size 50 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed, paraffin-embedded section of dog ischemic brain stained for Active/Cleaved CASP3 expression. Using CASP3 polyclonal antibody (Cat # PAB0243) at 1 : 2000. Staining is seen in the nuclei of dying neurons (black arrow) but not in the morphologically normal nuclei (open arrows) . CASP3 expression in the nucleus is considered to be a marker of Active/Cleaved CASP3 expression and apoptosis. Hematoxylin-eosin counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed, paraffin-embedded section of irradiated mouse spleen stained for Active/Cleaved CASP3 expression. Using CASP3 polyclonal antibody (Cat # PAB0243) at 1 : 2000. Staining is seen in the nuclei of a subset of the cell population. CASP3 expression in the nucleus is considered to be a marker of Active/Cleaved CASP3 expression and apoptosis. Hematoxylin-eosin counterstain.

Specification

Product Description	Rabbit polyclonal antibody raised against full length recombinant CASP3.
Immunogen	Recombinant protein corresponding to full length active human CASP3.
Host	Rabbit
Reactivity	Dog, Gerbils, Human, Mouse, Rat
Form	Liquid

Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In serum (0.05% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed, paraffin-embedded section of dog ischemic brain stained for Active/Cleaved CASP3 expression. Using CASP3 polyclonal antibody (Cat # PAB0243) at 1 : 2000. Staining is seen in the nuclei of dying neurons (black arrow) but not in the morphologically normal nuclei (open arrows) . CASP3 expression in the nucleus is considered to be a marker of Active/Cleaved CASP3 expression and apoptosis. Hematoxylin-eosin counterstain.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed, paraffin-embedded section of irradiated mouse spleen stained for Active/Cleaved CASP3 expression. Using CASP3 polyclonal antibody (Cat # PAB0243) at 1 : 2000. Staining is seen in the nuclei of a subset of the cell population. CASP3 expression in the nucleus is considered to be a marker of Active/Cleaved CASP3 expression and apoptosis. Hematoxylin-eosin counterstain.

- Immunoprecipitation

Gene Info — CASP3

Entrez GeneID	836
Gene Name	CASP3
Gene Alias	CPP32, CPP32B, SCA-1
Gene Description	caspase 3, apoptosis-related cysteine peptidase
Omim ID	600636
Gene Ontology	Hyperlink

Gene Summary

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq]

Other Designations

OTTHUMP00000165054|PARP cleavage protease|SREBP cleavage activity 1|Yama|apopain|caspase 3|caspase 3, apoptosis-related cysteine protease|cysteine protease CPP32|procaspase3

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [MAPK signaling pathway](#)
- [Natural killer cell mediated cytotoxicity](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)

Disease

- [Adenocarcinoma](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Clubfoot](#)
- [Colorectal Neoplasms](#)

- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung carcinoma](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [NARP](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Occupational Diseases](#)

- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Small Cell Lung Carcinoma](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)