

Goat Anti-Bovine IgG (H&L) secondary antibody (Peroxidase)

Catalog # PAB10432

Size 2 mL

Specification

Product Description	Goat anti-bovine IgG recognizes bovine IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.
Immunogen	Bovine IgG whole molecule
Host	Goat
Reactivity	Bovine
Form	Lyophilized
Conjugation	HRP
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 0.01 M Na ₂ HPO ₄ , 250 mM NaCl, pH 7.2.
Storage Instruction	Store vial at 4°C prior to restoration. For extended storage aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use.

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

- Dot Blot
- Enzyme Immunoassay

Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.