

Goat Anti-Horse IgG F(c) secondary antibody (Peroxidase)

Catalog # PAB10657

Size 2 mg

Specification

Product Description	Goat anti-horse IgG recognizes horse IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.
Immunogen	Horse IgG F(c) fragment
Host	Goat
Reactivity	Horse
Form	Lyophilized
Conjugation	HRP
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% gentamicin sulfate)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay
- Dot Blot

- Enzyme Immunoassay

Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.