## Chicken Anti-Rabbit IgG (H&L) secondary antibody (Rhodamine)

Catalog # PAB10808 Size 1 mg

Product Description Chicken anti-rabbit IgG recognizes rabbit IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Rhodamine.   Immunogen Rabbit IgG whole molecule   Host Chicken   Reactivity Rabbit   Form Lyophilized   Conjugation Rhodamine (TRITC)   Purification This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.   Conjugation Note 1.3 moles TRITC per mole of chicken IgG   Recommend Usage The optimal working dilution should be determined by the end user.   Storage Buffer Lyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))   Storage Instruction Store at 4°C c on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.   Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be delered freezing and thawing.	Specification	
HostChickenReactivityRabbitFormLyophilizedConjugationRhodamine (TRITC)PurificationThis product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.Conjugation Note1.3 moles TRITC per mole of chicken lgGRecommend UsageThe optimal working dilution should be determined by the end user.Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Product Description	
Reactivity Rabbit   Form Lyophilized   Conjugation Rhodamine (TRITC)   Purification This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.   Conjugation Note 1.3 moles TRITC per mole of chicken lgG   Recommend Usage The optimal working dilution should be determined by the end user.   Storage Buffer Lyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))   Storage Instruction Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.   Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Immunogen	Rabbit IgG whole molecule
FormLyophilizedConjugationRhodamine (TRITC)PurificationThis product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.Conjugation Note1.3 moles TRITC per mole of chicken lgGRecommend UsageThe optimal working dilution should be determined by the end user.Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should	Host	Chicken
ConjugationRhodamine (TRITC)PurificationThis product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.Conjugation Note1.3 moles TRITC per mole of chicken lgGRecommend UsageThe optimal working dilution should be determined by the end user.Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Reactivity	Rabbit
Purification This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.   Conjugation Note 1.3 moles TRITC per mole of chicken lgG   Recommend Usage The optimal working dilution should be determined by the end user.   Storage Buffer Lyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))   Storage Instruction Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.   Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Form	Lyophilized
Conjugation Note1.3 moles TRITC per mole of chicken lgGRecommend UsageThe optimal working dilution should be determined by the end user.Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Conjugation	Rhodamine (TRITC)
Recommend UsageThe optimal working dilution should be determined by the end user.Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Purification	
Storage BufferLyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))Storage InstructionStore at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.NoteThis product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Conjugation Note	1.3 moles TRITC per mole of chicken IgG
Storage Instruction Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.   Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Recommend Usage	The optimal working dilution should be determined by the end user.
After reconstitution with 1 mL of deionized water, store at -20°C or lower.   Aliquot to avoid repeated freezing and thawing.   Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul	Storage Buffer	
·	Storage Instruction	After reconstitution with 1 mL of deionized water, store at -20°C or lower.
	Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## Applications

- Immunofluorescence
- Flow Cytometry



## Publication Reference

• <u>Texas Red, a hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow</u> <u>microfluorometric and fluorescence microscopic studies.</u>

J A Titus, R Haugland, S O Sharrow, D M Segal.

Journal of Immunological Methods 1982 Apr; 50(2):193.