


TNNT2 polyclonal antibody


Catalog # PAB12744 Size 100 ug

Applications



Western Blot (Tissue lysate)

Tissue lysate from mouse heart was resolved onto 12% SDS-PAGE and transferred onto NC Membrane, then probed by TNNT2 polyclonal antibody (Cat # PAB12744) at 1 : 500, right lane. Left lane was depleted by immunizing peptide.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human cardiac muscle stained with TNNT2 polyclonal antibody (Cat # PAB12744) at 1 : 50 for 10 min at RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of TNNT2.
Immunogen	A synthetic peptide corresponding to internal domain of human TNNT2.
Host	Rabbit
Theoretical MW (kDa)	31
Reactivity	Cat, Dog, Human, Mouse, Rabbit
Specificity	This antibody recognizes 31 KDa of TNNT2 by Western blot.
Form	Liquid

Quality Control Testing

Antibody Reactive Against Synthetic Peptide.

Recommend Usage

Western Blot (0.1-1 ug/mL)
ELISA (0.01-0.1 ug/mL)
Immunoprecipitation (2-5 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer

In TBS, pH 7.2 (BSA, 10% Proclin300)

Storage Instruction

Store at 4°C. For long term storage store at -20°C or lower.
Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Tissue lysate)

Tissue lysate from mouse heart was resolved onto 12% SDS-PAGE and transferred onto NC Membrane, then probed by TNNT2 polyclonal antibody (Cat # PAB12744) at 1 : 500, right lane. Left lane was depleted by immunizing peptide.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human cardiac muscle stained with TNNT2 polyclonal antibody (Cat # PAB12744) at 1 : 50 for 10 min at RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

- Immunoprecipitation

- Enzyme-linked Immunoabsorbent Assay

Gene Info — TNNT2

Entrez GeneID[7139](#)**Gene Name**

TNNT2

Gene Alias

CMH2, CMPD2, MGC3889, RCM3, TnTC, cTnT

Gene Description

troponin T type 2 (cardiac)

Omim ID[115195 191045 601494](#)**Gene Ontology**[Hyperlink](#)

Gene Summary

The protein encoded by this gene is the tropomyosin-binding subunit of the troponin complex, which is located on the thin filament of striated muscles and regulates muscle contraction in response to alterations in intracellular calcium ion concentration. Mutations in this gene have been associated with familial hypertrophic cardiomyopathy as well as with dilated cardiomyopathy. Transcripts for this gene undergo alternative splicing that results in many tissue-specific isoforms, however, the full-length nature of some of these variants has not yet been determined. [provided by RefSeq]

Other Designations

OTTHUMP00000033864|OTTHUMP00000033867|cardiomyopathy, dilated 1D (autosomal dominant)|cardiomyopathy, hypertrophic 2|troponin T type 2, cardiac|troponin T, cardiac muscle|troponin T2, cardiac

Pathway

- [Cardiac muscle contraction](#)
- [Hypertrophic cardiomyopathy \(HCM\)](#)

Disease

- [Cardiomegaly](#)
- [Cardiomyopathies](#)
- [Cardiomyopathy](#)
- [Cardiovascular Diseases](#)
- [Death](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hypertrophy](#)
- [Lipid Metabolism](#)
- [Mitochondrial Myopathies](#)
- [Tachycardia](#)