


IL2RB polyclonal antibody

Catalog # PAB18399 Size 100 ug


Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from COS-7 cells, using IL2RB polyclonal antibody (Cat # PAB18399).

Peptide "+" means "peptide blocking".



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using IL2RB polyclonal antibody (Cat # PAB18399).

Peptide "+" means "peptide blocking".

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of IL2RB.
Immunogen	A synthetic peptide corresponding to residues surrounding Y364 of human IL2RB.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Specificity	This antibody is specific to IL2RB.
Form	Liquid

Purification	Affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:100) ELISA (1:20000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of extracts from COS-7 cells, using IL2RB polyclonal antibody (Cat # PAB18399).
Peptide "+" means "peptide blocking".

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using IL2RB polyclonal antibody (Cat # PAB18399).
Peptide "+" means "peptide blocking".

- Enzyme-linked Immunoabsorbent Assay

Gene Info — IL2RB

Entrez GeneID	3560
Protein Accession#	P14784
Gene Name	IL2RB
Gene Alias	CD122, P70-75
Gene Description	interleukin 2 receptor, beta
Omim ID	146710
Gene Ontology	Hyperlink

Gene Summary

The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is present in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha subunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha/beta subunit heterodimer, while the high affinity form consists of an alpha/beta/gamma subunit heterotrimer. Both the intermediate and high affinity forms of the receptor are involved in receptor-mediated endocytosis and transduction of mitogenic signals from interleukin 2. The protein encoded by this gene represents the beta subunit and is a type I membrane protein. [provided by RefSeq]

Other Designations

CD122 antigen|OTTHUMP00000028799|high affinity IL-2 receptor beta subunit|interleukin 2 receptor beta

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bipolar Disorder](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Chagas Disease](#)
- [Colitis](#)
- [Connective Tissue Diseases](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)

- [Disease Progression](#)
- [Edema](#)
- [Endometriosis](#)
- [Esophagitis](#)
- [Fetal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Hyperparathyroidism](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leishmaniasis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Measles](#)
- [Multiple Sclerosis](#)
- [Mumps](#)
- [Musculoskeletal Diseases](#)
- [Myasthenia Gravis](#)
- [Occupational Diseases](#)
- [Pain](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Prostate cancer](#)

- [Prostatic Neoplasms](#)
- [Radiation Injuries](#)
- [Radiation Pneumonitis](#)
- [Respiratory Syncytial Virus Infections](#)
- [Rubella](#)
- [Schizophrenia](#)
- [Skin Diseases](#)
- [Tobacco Use Disorder](#)
- [Viremia](#)
- [Wegener Granulomatosis](#)