


NFKBIB polyclonal antibody

Catalog # PAB18763

Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of HepG2 cell lysate with NFKBIB polyclonal antibody (Cat # PAB18763) at 1: 200 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of NFKBIB.
Immunogen	A synthetic peptide corresponding to 14 amino acids at C-terminus of human NFKBIB.
Host	Rabbit
Reactivity	Human
Form	Liquid
Recommend Usage	ELISA (1:20000-1:80000) Western Blot (1:200-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In buffer containing 0.02% sodium azide
Storage Instruction	Store at 4°C for three months. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of HepG2 cell lysate with NFKBIB polyclonal antibody (Cat # PAB18763) at 1: 200 dilution.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — NFKBIB

Entrez GeneID [4793](#)

Protein Accession# [NM_002503](#)

Gene Name NFKBIB

Gene Alias IKBB, TRIP9

Gene Description nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, beta

Omim ID [604495](#)

Gene Ontology [Hyperlink](#)

Gene Summary NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM 164014), or RELB (MIM 604758) to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664 or IKBKB, MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]

Other Designations -

Pathway

- [Adipocytokine signaling pathway](#)
- [B cell receptor signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [T cell receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Atherosclerosis](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Connective Tissue Diseases](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Empyema](#)
- [Fetal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [HIV Infections](#)
- [Inflammation](#)
- [Lymphoma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pneumococcal Infections](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)

- [Skin Diseases](#)