
# WDR25 polyclonal antibody


Catalog # PAB20165 Size 100 uL

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human testis with WDR25 polyclonal antibody (Cat # PAB20165) shows strong nuclear positivity in bone marrow poietic cells at 1:50-1:200 dilution.




### Western Blot

Western blot analysis of Lane 1: RT-4, Lane 2: U-251 MG, Lane 3: Human Plasma, Lane 4: Liver, Lane 5: Tonsil with WDR25 polyclonal antibody (Cat # PAB20165) at 1:250-1:500 dilution.

## Specification

| | |
|---|---|
| **Product Description** | Rabbit polyclonal antibody raised against recombinant WDR25. |
| **Immunogen** | Recombinant protein corresponding to amino acids of human WDR25. |
| **Sequence** | LVAYDDSDSEAETEHAGSFNATGQQKDTSGVARPPGQDFASGTLDVPKAGAQPTKHGSCEDP GGYRLPLAQLGRSDRGSCPSQRLQWPGKEPQVTFPIKEPSCSSLWTS |
| **Host** | Rabbit |
| **Reactivity** | Human |
| **Form** | Liquid |

Copyright © 2023 Abnova Corporation. All Rights Reserved.

| | |
|---|---|
| **Purification** | Antigen affinity purification |
| **Isotype** | IgG |
| **Recommend Usage** | Immunohistochemistry (1:50-1:200)<br>Western Blot (1:250-1:500)<br>The optimal working dilution should be determined by the end user. |
| **Storage Buffer** | In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide) |
| **Storage Instruction** | Store at 4°C. For long term storage store at -20°C.<br>Aliquot to avoid repeated freezing and thawing. |
| **Note** | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only. |

## Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining of human testis with WDR25 polyclonal antibody (Cat # PAB20165) shows strong nuclear positivity in bone marrow poietic cells at 1:50-1:200 dilution.

- Western Blot

  Western blot analysis of Lane 1: RT-4, Lane 2: U-251 MG, Lane 3: Human Plasma, Lane 4: Liver, Lane 5: Tonsil with WDR25 polyclonal antibody (Cat # PAB20165) at 1:250-1:500 dilution.

## Gene Info — WDR25

| | |
|---|---|
| **Entrez GeneID** | 79446 |
| **Protein Accession#** | Q64LD2 |
| **Gene Name** | WDR25 |
| **Gene Alias** | MGC4645 |
| **Gene Description** | WD repeat domain 25 |
| **Gene Ontology** | Hyperlink |
| **Other Designations** | pre-mRNA splicing factor-like |

Copyright © 2023 Abnova Corporation. All Rights Reserved.