


MYC (phospho S62) polyclonal antibody

Catalog # PAB25882 Size 100 ug

Applications



Immunofluorescence

Immunofluorescence staining of methanol-fixed HeLa cells using MYC (phospho S62) polyclonal antibody (Cat # PAB25882, red).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of MYC.
Immunogen	Synthetic phosphopeptide corresponding to residues surrounding S62 of human MYC.
Sequence	P-L-Sp-P-S
Host	Rabbit
Theoretical MW (kDa)	60
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Affinity chromatography
Concentration	1 mg/mL
Recommend Usage	Immunofluorescence (1:100-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (without Mg^{2+} and Ca^{2+}), 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence

Immunofluorescence staining of methanol-fixed HeLa cells using MYC (phospho S62) polyclonal antibody (Cat # PAB25882, red).

Gene Info — MYC

Entrez GeneID[4609](#)**Protein Accession#**[P01106](#)**Gene Name**

MYC

Gene Alias

bHLHe39, c-Myc

Gene Description

v-myc myelocytomatosis viral oncogene homolog (avian)

Omim ID[113970 190080](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemia and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq]

Other Designations

avian myelocytomatosis viral oncogene homolog|myc proto-oncogene protein|v-myc avian myelocytomatosis viral oncogene homolog

Pathway

- [Acute myeloid leukemia](#)

- [Bladder cancer](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [TGF-beta signaling pathway](#)
- [Thyroid cancer](#)
- [Wnt signaling pathway](#)

Disease

- [Alzheimer disease](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chromosome Aberrations](#)
- [Colorectal Neoplasms](#)
- [Diabetes Complications](#)
- [Disease Progression](#)
- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)

- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Mouth Neoplasms](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Thyroid Diseases](#)
- [Thyroid Neoplasms](#)
- [Translocation](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)