


XRCC6 polyclonal antibody


Catalog # PAB27138 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of HeLa cell lysate with XRCC6 polyclonal antibody (Cat # PAB27138).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue using XRCC6 polyclonal antibody (Cat # PAB27138).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of XRCC6.
Immunogen	A synthetic peptide corresponding to human XRCC6.
Host	Rabbit
Theoretical MW (kDa)	70
Reactivity	Human, Mouse
Specificity	XRCC6 polyclonal antibody detects endogenous levels of XRCC6 protein.
Form	Liquid

Purification	Antigen affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (0.05% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of HeLa cell lysate with XRCC6 polyclonal antibody (Cat # PAB27138).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue using XRCC6 polyclonal antibody (Cat # PAB27138).

Gene Info — XRCC6

Entrez GeneID	2547
Protein Accession#	P12956
Gene Name	XRCC6
Gene Alias	CTC75, CTCBF, G22P1, KU70, ML8, TLAA
Gene Description	X-ray repair complementing defective repair in Chinese hamster cells 6
Omim ID	152690
Gene Ontology	Hyperlink

Gene Summary

The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq]

Other Designations

ATP-dependent DNA helicase II, 70 kDa subunit|CTC box binding factor 75 kDa subunit|Ku autoantigen p70 subunit|Ku autoantigen, 70kDa|OTTHUMP00000028581|thyroid autoantigen 70kD (Ku antigen)|thyroid autoantigen 70kDa (Ku antigen)|thyroid-lupus autoantigen

Pathway

- [Non-homologous end-joining](#)

Disease

- [Adenocarcinoma](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Central Nervous System Neoplasms](#)
- [Diabetes Mellitus](#)
- [DNA Damage](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)

- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Male Urogenital Diseases](#)
- [Micronuclei](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pterygium](#)
- [Radiodermatitis](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)