


BID polyclonal antibody

Catalog # PAB27139

Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of Jurkat cell lysate with treatment of Etoposide 25uM 60min treated. Using BID polyclonal antibody (Cat # PAB27139).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human brain tissue using BID polyclonal antibody (Cat # PAB27139).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of BID.
Immunogen	A synthetic peptide corresponding to human BID.
Host	Rabbit
Theoretical MW (kDa)	25
Reactivity	Human, Mouse
Specificity	BID polyclonal antibody detects endogenous levels of BID protein.
Form	Liquid

Purification	Antigen affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (0.05% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of Jurkat cell lysate with treatment of Etoposide 25uM 60min treated. Using BID polyclonal antibody (Cat # PAB27139).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human brain tissue using BID polyclonal antibody (Cat # PAB27139).

Gene Info — BID

Entrez GeneID	637
Protein Accession#	P55957
Gene Name	BID
Gene Alias	FP497, MGC15319, MGC42355
Gene Description	BH3 interacting domain death agonist
Omim ID	601997
Gene Ontology	Hyperlink

Gene Summary

This gene encodes a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but the full-length nature of some variants has not been defined. [provided by RefSeq]

Other Designations

BH3-interacting domain death agonist|BID isoform ES(1b)|BID isoform L(2)|BID isoform Si6|Human BID coding sequence|OTTHUMP00000196197|apoptotic death agonist|desmocollin type 4

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Natural killer cell mediated cytotoxicity](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)

Disease

- [Adenocarcinoma](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Clubfoot](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Disease Progression](#)
- [Edema](#)

- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Kidney Failure](#)
- [Lymphatic Metastasis](#)
- [Neoplasm Metastasis](#)
- [Stomach Neoplasms](#)
- [Tobacco Use Disorder](#)