


CD79B polyclonal antibody


Catalog # PAB29311 Size 100 uL

Applications




Western Blot (Transfected lysate)

Western blot analysis of Lane 1: Negative control (vector only transfected HEK293T lysate), Lane 2: Over-expression lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells) with PIK3R1 polyclonal antibody (Cat# PAB29311) at 1:100 - 1:250 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lymph node tissue with CD79B polyclonal antibody (Cat# PAB29311) at 1:200 - 1:500 dilution.



Immunofluorescence

Immunofluorescence staining of U-2 OS cells with CD79B polyclonal antibody (Cat# PAB29311) under 1-4 ug/mL working concentration.

Specification

Product Description	Rabbit polyclonal antibody raised against recombinant human CD79B.
Immunogen	Recombinant protein corresponding to amino acids 182-229 of human CD79B.
Sequence	DKDDSKAGMEEDHTYEGLDIDQTATYEDIVTLRTGEVKWSVGEHPGQE
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunofluorescence (1-4 ug/mL) Immunohistochemistry (1:200-1:500) Western Blot (1:100-1:250) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western blot analysis of Lane 1: Negative control (vector only transfected HEK293T lysate), Lane 2: Over-expression lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells) with PIK3R1 polyclonal antibody (Cat# PAB29311) at 1:100 - 1:250 dilution.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lymph node tissue with CD79B polyclonal antibody (Cat# PAB29311) at 1:200 - 1:500 dilution.

- Immunofluorescence

Immunofluorescence staining of U-2 OS cells with CD79B polyclonal antibody (Cat# PAB29311) under 1-4 ug/mL working concentration.

Gene Info — CD79B

Entrez GeneID	974
Protein Accession#	P40259
Gene Name	CD79B
Gene Alias	B29, IGB
Gene Description	CD79b molecule, immunoglobulin-associated beta
Omim ID	147245
Gene Ontology	Hyperlink
Gene Summary	The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-beta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq]
Other Designations	B-cell antigen receptor complex-associated protein beta chain B-cell-specific glycoprotein B29 CD79B antigen CD79b antigen (immunoglobulin-associated beta) Ig-beta immunoglobulin-associated B29

Pathway

- [B cell receptor signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Lymphoproliferative Disorders](#)

- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)